

COMBINATORIAL ASPECTS OF MIRROR SYMMETRY

VICTOR BATYREV AND BENJAMIN NILL

ABSTRACT. The purpose of this paper is to review some combinatorial ideas behind the mirror symmetry for Calabi-Yau hypersurfaces and complete intersections in Gorenstein toric Fano varieties. We suggest as a basic combinatorial object the notion of a Gorenstein polytope of index r . A natural combinatorial duality for d -dimensional Gorenstein polytopes of index r extends the well-known polar duality for reflexive polytopes (case $r = 1$). We consider the Borisov duality between two nef-partitions as a duality between two Gorenstein polytopes P and P^* of index r together with selected special $(r - 1)$ -dimensional simplices $S \subset P$ and $S' \subset P^*$. Different choices of these simplices suggest an interesting relation to Homological Mirror Symmetry.

INTRODUCTION

Several papers of the first author and Borisov were devoted to a combinatorial method for explicit constructions of mirror pairs of Calabi-Yau varieties (see [Bor93, Bat94, BB96a, BB97]). A starting point of this method was the notion of a *reflexive polytope* introduced in [Bat94]. A reflexive polytope $P \subset \mathbb{R}^d$ is a convex d -dimensional polytope with vertices in $\mathbb{Z}^n$ containing the origin $0 \in \mathbb{R}^d$ in its interior such that its dual (polar) polytope

$$P^* := \{y = (y_1, \dots, y_d) \in \mathbb{R}^d : \sum_{k=1}^d x_k y_k \geq -1 \ \forall x = (x_1, \dots, x_d) \in P\}.$$

is again a lattice polytope (i.e., all vertices of P^* belong to $\mathbb{Z}^n$). In this case P^* is also a reflexive polytope and $(P^*)^* = P$. For any convex lattice polytope $\Delta \subset \mathbb{R}^d$ we denote by X_Δ a generic hypersurface in $(\mathbb{C}^*)^d$ defined by the equation

$$F_\Delta(z) := \sum_{m \in \mathbb{Z}^d \cap \Delta} a_m z^m = 0,$$

where the coefficients a_m ($m \in \mathbb{Z}^d \cap P$) are independent variables and $z = (z_1, \dots, z_d)$ are standard complex coordinates on $(\mathbb{C}^*)^d$. If P is reflexive, then $X_P \subset (\mathbb{C}^*)^d$ admits a natural compactification $\overline{X}_P$ which is a $(d - 1)$ -dimensional Calabi-Yau variety. Moreover, the duality between P and P^* becomes the mirror duality between $(d - 1)$ -dimensional Calabi-Yau varieties $\overline{X}_P$ and $\overline{X}_{P^*}$.

Many explicit constructions of mirrors of Calabi-Yau complete intersections suggested by the first author and van Straten in [BS95] motivated Borisov to generalize the polar duality for reflexive polytopes to a more general duality for so called *nef-partitions* (see [Bor93]). From a combinatorial

point of view, a *nef-partition of length r* is a decomposition $P = P_1 + \cdots + P_r$ of a d -dimensional reflexive polytope $P \subset \mathbb{R}^n$ into a Minkowski sum of r lattice polytopes $P_1, \dots, P_r$ such that $0 \in P_i$ for all $i = 1, \dots, r$ (it is not assumed that each P_i has the maximal dimension d). It can be shown that for any $j = 1, \dots, r$ the polytope

$$Q_j = \{y \in \mathbb{R}^d : \sum_{k=1}^d x_k y_k \geq -\delta_{ij} \quad \forall x \in P_i, \quad \forall i = 1, \dots, r\}$$

has vertices in $\mathbb{Z}^d$, $0 \in Q_j$ and the Minkowski sum $Q = Q_1 + \cdots + Q_r$ is again a d -dimensional reflexive polytope. Moreover, the lattice polytopes $P_1, \dots, P_r$ can be obtained from lattice polytopes $Q_1, \dots, Q_r$ by the same formula:

$$P_i = \{y \in \mathbb{R}^d : \sum_{k=1}^d x_k y_k \geq -\delta_{ji} \quad \forall x \in Q_j, \quad \forall j = 1, \dots, r\}.$$

The Minkowski sum decomposition $Q = Q_1 + \cdots + Q_r$ is called the *dual nef-partition*. Some examples of nef-partitions and their duals are given in Examples 3.2, 3.14 and 3.15. If $X_{\{P_i\}}$ is a generic complete intersection in $(\mathbb{C}^*)^d$ defined by generic equations

$$F_{P_1}(z) = \cdots = F_{P_r}(z) = 0,$$

then $X_{\{P_i\}}$ admits a natural compactification $\overline{X}_{\{P_i\}}$ which is a $(d-r)$ -dimensional Calabi-Yau variety. Moreover, the duality of nef-partitions $\{P_i\}$ and $\{Q_j\}$ becomes the mirror duality between $(d-r)$ -dimensional Calabi-Yau varieties $\overline{X}_{\{P_i\}}$ and $\overline{X}_{\{Q_j\}}$.

If $P \subset \mathbb{R}^n$ is a reflexive polytope, $P = P_1 + \cdots + P_r$ is a nef-partition and $Q = Q_1 + \cdots + Q_r$ is the dual nef-partition, then the lattice polytopes $P_1, \dots, P_r$ satisfy some additional combinatorial conditions:

1. $\mathbb{R}_{\geq 0}(P_k) \cap \mathbb{R}_{\geq 0}(P_l) = 0$ for all $1 \leq k < l \leq r$;
2. the convex hull $\text{Conv}\{P_1, \dots, P_r\}$ is the dual reflexive polytope Q^* ;
3. $Q^* \cap \mathbb{Z}^d = (P_1 \cap \mathbb{Z}^d) \cup \cdots \cup (P_r \cap \mathbb{Z}^d)$. Moreover, $m \in Q^* \cap \mathbb{Z}^d$ is a vertex of Q^* if and only if m is a nonzero vertex of P_i for some $i \in \{1, \dots, r\}$.

For lattice polytopes $\Delta_1, \dots, \Delta_r \in \mathbb{R}^d$ we denote by $\Delta_1 * \cdots * \Delta_r$ the polytope in $\mathbb{R}^d \times \mathbb{R}^r$ which is the convex hull

$$\text{Conv}\{(\Delta_1 \times e_1), \dots, (\Delta_r \times e_r)\},$$

where $e_1, \dots, e_r$ is the standard basis of $\mathbb{R}^r$. We call $\Delta_1 * \cdots * \Delta_r$ *Cayley polytope associated with $\Delta_1, \dots, \Delta_r$* .

Let P and Q be two d -dimensional reflexive polytopes and $P = P_1 + \cdots + P_r$ and $Q = Q_1 + \cdots + Q_r$ nef-partitions which are dual to each other. We consider two $(d+r-1)$ -dimensional polytopes $\tilde{P} := P_1 * \cdots * P_r$, $\tilde{Q} := Q_1 * \cdots * Q_r$ and two $(d+r)$ -dimensional cones

$$C_{\tilde{P}} := \mathbb{R}_{\geq 0}(P_1 \times e_1) + \cdots + \mathbb{R}_{\geq 0}(P_r \times e_r) \subset \mathbb{R}^d \times \mathbb{R}^r,$$

$$C_{\tilde{Q}} := \mathbb{R}_{\geq 0}(Q_1 \times e_1) + \cdots + \mathbb{R}_{\geq 0}(Q_r \times e_r) \subset \mathbb{R}^d \times \mathbb{R}^r.$$

It was proved in [BB97, Theorem 4.6] that the cones $C_{\tilde{P}}$ and $C_{\tilde{Q}}$ are dual to each other with respect to standard scalar product on $\mathbb{R}^d \times \mathbb{R}^r$. Moreover, there exists a unique lattice point $p \in r\tilde{P}$ (resp. $q \in r\tilde{Q}$) in the relative interior of $r\tilde{P}$ (resp. $r\tilde{Q}$) such that $r\tilde{P} - p$ (resp. $r\tilde{Q} - q$) is a $(d + r - 1)$ -dimensional reflexive polytope. For this reason, the duality for nef-partitions of length r can be considered as a special case of the more general duality for so called *reflexive Gorenstein cones of index r* . The duality for reflexive Gorenstein cones gives new possibilities for constructing mirrors of Calabi-Yau varieties. In particular, it allows to construct mirrors of rigid-Calabi-Yau 3-folds in form of so called *generalized Calabi-Yau varieties* or *Landau-Ginzburg orbifolds* [CDP93, BB97].

Recall that a $(d + r - 1)$ -dimensional lattice polytope $P \subset \mathbb{R}^{d+r-1}$ is called a *Gorenstein polytope of index r* , if rP contains a single interior lattice point p and $rP - p$ is a reflexive polytope. In this case, the $(d + r)$ -dimensional cone

$$C_P := \mathbb{R}_{\geq 0}(P \times 1) \subset \mathbb{R}^{d+r-1} \times \mathbb{R}.$$

is called *reflexive Gorenstein cone of index r* associated with P . Let

$$C_P^\vee := \{y \in \mathbb{R}^d \times \mathbb{R}, \sum_k x_k y_k \geq 0, \forall x \in C_P\}$$

be the dual cone. Then there exists another Gorenstein polytope P^* of index r such that C_{P^*} is isomorphic (via the action of $GL(d + r, \mathbb{Z})$) to the cone $C_P^\vee$. The polytope P^* is uniquely determined up to the action of the affine group $AGL(d + r - 1, \mathbb{Z})$ on $\mathbb{R}^{d+r-1}$. It is called the *dual Gorenstein polytope of index r* . In this paper, we want to look at the duality of reflexive Gorenstein cones of index r from the viewpoint of the duality of Gorenstein polytopes of index r . For $r = 1$ the latter coincides with the already known polar duality for reflexive polytopes. It is important to consider an additional structure on Gorenstein polytopes P which is defined by a choice of a special $(r - 1)$ -dimensional lattice simplex $S \subset P$. The combinatorial notion of a *special simplex* was recently introduced in [Ath05] in order to prove a conjecture of Stanley. If a Gorenstein polytope P of index r and its dual P^* both contain special $(r - 1)$ -simplices, we naturally obtain two nef-partitions dual to each other. Moreover, the existence of special simplices in P and P^* can be considered as another characterization of nef-partitions and their duals (see Proposition 3.6). The choice of special $(r - 1)$ -simplices is not unique. Therefore, one may obtain many different nef-partitions from the same pair of Gorenstein polytopes (P, P^*) . We expect that different choices of special $(r - 1)$ -simplices $S \subset P$ and $S' \subset P^*$ define Calabi-Yau complete intersections which are equivalent from the viewpoint of Homological Mirror Symmetry.

The paper is organized as follows. In Section 1 we recall the notions of Gorenstein polytopes, reflexive Gorenstein cones and their dualities. In

Section 2 we establish a bijective correspondence between Cayley structures of length r on a Gorenstein polytope P of index r and special $(r - 1)$ -simplices in the dual Gorenstein polytope P^* . In Section 3 we review the notion of nef-partitions and their duality as well as combinatorial properties of nef-partitions. In Section 4 we define the E_{st} -function of a Gorenstein polytope P and formulate some open questions and conjectures. Finally, in Section 5 we give a precise formulation of a conjecture which connects special $(r - 1)$ -simplices in a Gorenstein polytope P with the Homological Mirror Symmetry. In the last Section 6 we consider as an addendum to Section 3 some combinatorial operations with nef-partitions.

Acknowledgment. We are grateful to anonymous referees for valuable remarks. The second author thanks Christian Haase for discussions, motivating questions and suggestions.

1. GORENSTEIN POLYTOPES AND REFLEXIVE GORENSTEIN CONES

1.1. Gorenstein polytopes. Let $M \cong \mathbb{Z}^d$ be a lattice of rank d , $N = \text{Hom}_{\mathbb{Z}}(M, \mathbb{Z})$ the dual lattice, $\langle \cdot, \cdot \rangle : M \times N \rightarrow \mathbb{Z}$ the natural pairing. We define $M_{\mathbb{Q}} := M \otimes_{\mathbb{Z}} \mathbb{Q}$, $M_{\mathbb{R}} := M \otimes_{\mathbb{Z}} \mathbb{R}$, $N_{\mathbb{Q}} := N \otimes_{\mathbb{Z}} \mathbb{Q}$, $N_{\mathbb{R}} := N \otimes_{\mathbb{Z}} \mathbb{R}$. By a polytope $\Delta \subset M_{\mathbb{R}}$ we always mean a convex hull of finitely many points in $M_{\mathbb{R}}$. The set of vertices of Δ is denoted by $\mathcal{V}(\Delta)$, the relative interior by $\text{int}(\Delta)$, and the boundary by $\partial\Delta$. For arbitrary subset $A \subseteq M_{\mathbb{R}}$ we denote by $\text{Conv}(A)$ (resp. $\text{aff}(A)$) the convex (resp. affine) hull of A .

From now on we assume $\Delta \subseteq M_{\mathbb{R}}$ to be a d -dimensional *lattice polytope*, i.e., $\mathcal{V}(\Delta) \subset M$.

Definition 1.1. Assume that Δ has the origin $0 \in M$ in its interior. Then the *dual polytope* of Δ is defined as

$$\Delta^* := \{y \in N_{\mathbb{R}} : \langle y, x \rangle \geq -1 \forall x \in \Delta\}.$$

This is a rational polytope (i.e., $\mathcal{V}(\Delta^*) \subseteq N_{\mathbb{Q}}$) containing the origin $0 \in N$ in its interior.

If Δ contains an interior lattice point $m \in M$ (not necessary 0), we say that $(\Delta - m)^*$ is the *dual polytope of Δ with respect to m* .

Definition 1.2. Let $\Delta \subseteq M_{\mathbb{R}}$ be a lattice polytope containing the origin $0 \in M$ in its interior. We say Δ is *reflexive*, if the dual polytope Δ^* is also a lattice polytope. In this case, $0 \in \Delta$ is a single interior lattice point.

In general, if a lattice polytope $\Delta \subset M_{\mathbb{R}}$ contains a single interior lattice point m (not necessary 0), we say that Δ is *reflexive with respect to m* , if $\Delta - m$ is reflexive.

Remark 1.3. It is easy to see that a lattice polytope $\Delta \subseteq M_{\mathbb{R}}$ is reflexive if and only if every facet of Δ has integral distance 1 from its unique interior lattice point. Recall that a point $x \in M_{\mathbb{R}}$ is said to have *integral distance* δ from a facet F of Δ , if there is a primitive lattice point $u \in N$ such that $u \perp F$ and $\langle u, F \rangle - \langle u, x \rangle = \delta$.

Remark 1.4. The following example shows that in dimension ≥ 3 there exist lattice polytopes containing 0 as a single interior lattice point which are not reflexive. Let Δ be the 3-dimensional lattice polytope with vertices $\pm(1, 0, 0)$, $\pm(0, 1, 0)$, $\pm(1, 1, 2)$. Then $(0, 0, 0) \in \Delta$ is a single interior lattice point, however, the integral distance from 0 to the facet

$$F := \text{Conv}((1, 0, 0), (0, 1, 0), (1, 1, 2))$$

is 2, since for the primitive vector $u := (2, 2, -1)^t$ we have $u \perp F$ and $\langle u, F \rangle = 2$.

Definition 1.5. Let r be a positive integer. A lattice d -dimensional polytope $\Delta \subset M_{\mathbb{R}}$ is called *Gorenstein of index r* , if $r\Delta$ contains an interior lattice point m and $r\Delta - m$ is reflexive.

Example 1.6. Let $\Delta \subset \mathbb{R}^d$ be the standard unit cube: $0 \leq x_i \leq 1 \ \forall i = 1, \dots, d$. Then $2\Delta - (1, \dots, 1)$ is reflexive. Therefore Δ is a Gorenstein polytope of index 2.

It is clear that Gorenstein polytopes can be characterized as follows:

Remark 1.7. A lattice polytope Δ is a Gorenstein polytope of index r if and only if there is a rational point $x \in \text{int}(\Delta) \cap \frac{1}{r}M$ having the integral distance $1/r$ from any facet of Δ .

1.2. Reflexive Gorenstein cones. Let $\overline{M}$ and $\overline{N}$ be lattices of rank $\overline{d}$ which are dual to each other. Let $\sigma \subset \overline{M}_{\mathbb{R}}$ be a $\overline{d}$ -dimensional rational finite polyhedral cone with vertex $0 \in \overline{M}$. Then

$$\sigma^{\vee} := \{y \in \overline{N}_{\mathbb{R}} : \langle x, y \rangle \geq 0 \ \forall x \in \sigma\}$$

is called the *dual cone* of σ . It is also a $\overline{d}$ -dimensional rational finite polyhedral cone with vertex 0.

We recall some definitions from [BB97]:

Definition 1.8. A $\overline{d}$ -dimensional rational finite polyhedral cone σ is called *Gorenstein cone*, if it is generated by finitely many lattice points which are contained in an affine hyperplane $\{x \in \overline{M}_{\mathbb{R}} : \langle x, n \rangle = 1\}$ for some $n \in \overline{N}$. In this case, the lattice point $n \in \text{int}(\sigma^{\vee})$ is uniquely determined, and we denote it by n_{σ} . One has $\text{int}(\sigma^{\vee}) \cap \overline{N} = n_{\sigma} + \sigma^{\vee} \cap \overline{N}$. We define the k th slice of σ as $\sigma_{(k)} := \sigma \cap \{x \in \overline{M}_{\mathbb{R}} : \langle x, n_{\sigma} \rangle = k\}$.

Definition 1.9. Let $\sigma \subseteq \overline{M}_{\mathbb{R}}$ be a Gorenstein cone. Then the lattice polytope $\tilde{\Delta} := \sigma_{(1)}$ is called the *support* of σ ; it is a $(\overline{d} - 1)$ -dimensional lattice polytope with respect to the affine lattice $\widetilde{M} := \overline{M} \cap \{x \in \overline{M}_{\mathbb{R}} : \langle x, n_{\sigma} \rangle = 1\}$.

Let $\Delta \subseteq M_{\mathbb{R}}$ be a lattice polytope of dimension d . We set $\overline{M} := M \oplus \mathbb{Z}$, $\overline{d} := d + 1$, $\widetilde{M} := M \times \{1\}$, $\tilde{\Delta} := \Delta \times \{1\}$. Then $\mathbb{R}_{\geq 0}(\tilde{\Delta})$ is a Gorenstein cone with support $\tilde{\Delta}$.

Using the two constructions above, we obtain a 1-to-1 correspondence between $\bar{d}$ -dimensional Gorenstein cones $\sigma \subset \overline{M}$ considered up to action of $GL(\bar{d}, \mathbb{Z})$ on $\overline{M}$ and $(\bar{d}-1)$ -dimensional lattice polytopes $\tilde{\Delta} \subset \overline{M}_{\mathbb{R}}$ considered up to an affine transformation from $AGL(\bar{d}-1, \mathbb{Z})$.

Definition 1.10. A Gorenstein cone σ is called *reflexive*, if $\sigma^{\vee}$ is also a Gorenstein cone. In this case, we denote by $m_{\sigma^{\vee}}$ the unique lattice point in $\overline{M}$ such that the affine hyperplane $\{y \in \overline{N}_{\mathbb{R}} : \langle m_{\sigma^{\vee}}, y \rangle = 1\}$ contains all lattice points generating the cone $\sigma^{\vee}$. The number

$$r = \langle m_{\sigma^{\vee}}, n_{\sigma} \rangle$$

is called the *index* of the reflexive Gorenstein cone σ . Using $(\sigma^{\vee})^{\vee} = \sigma$, we see that r is also the index of the reflexive Gorenstein cone $\sigma^{\vee}$. Thus, we obtain a natural duality of reflexive Gorenstein cones of index r .

It was observed in [BB97, Prop.2.11] that reflexive Gorenstein cones correspond to Gorenstein polytopes:

Proposition 1.11. *Let σ be a $\bar{d}$ -dimensional Gorenstein cone with support $\tilde{\Delta}$. Then the following statements are equivalent:*

- (1) σ is a reflexive Gorenstein cone of index r ;
- (2) $\sigma_{(r)}$ is a reflexive polytope;
- (3) $\tilde{\Delta}$ is a Gorenstein polytope of index r

In this case, $\sigma_{(r)}$ (resp. $(\sigma^{\vee})_{(r)}$) is a reflexive polytope with the unique interior lattice point $m_{\sigma^{\vee}}$ (resp. n_{σ}).

The duality of reflexive Gorenstein cones of index r defines a natural duality of Gorenstein polytopes of index r :

Definition 1.12. Let $\Delta \subseteq M_{\mathbb{R}}$ be a d -dimensional Gorenstein polytope of index r . Then the *dual Gorenstein polytope* Δ^* is defined as the support of the dual cone $\sigma^{\vee}$, where σ is the reflexive Gorenstein cone associated to Δ . By 1.11, Δ^* is also a d -dimensional Gorenstein polytope of index r .

Remark 1.13. We want to compare the duality of d -dimensional Gorenstein polytopes of index r with the polar duality of reflexive polytopes. If $r = 1$, then the dualities are the same. For any $r \geq 1$ two Gorenstein polytopes Δ and Δ^* are combinatorially dual to each other. However, if $r > 1$, then the two reflexive polytopes $r\Delta$ and $r\Delta^*$ are *not* dual to each other as reflexive polytopes. In order to obtain the dual reflexive polytope $(r\Delta)^*$ from the Gorenstein polytope Δ^* one must replace the lattice N by another one (see 1.14 and 1.15 below).

Let us consider the duality of Gorenstein polytopes of index r in more detail.

1.3. Duality of Gorenstein polytopes.

Definition 1.14. Let $\sigma \subseteq \overline{M}_{\mathbb{R}}$ be a Gorenstein cone of index r with support $\tilde{\Delta} \subseteq \widetilde{M}_{\mathbb{R}}$. We define two lattices

$$\overline{M}_{(r)} := \{x \in \overline{M} : \langle x, n_{\sigma} \rangle = 0 \pmod{r}\}, \text{ and } \overline{N}^{(r)} := \overline{N} + \frac{1}{r}\mathbb{Z}n_{\sigma}.$$

The lattice $\overline{M}_{(r)}$ is dual to $\overline{N}^{(r)}$ and one has $\overline{M}_{(r)} \subseteq \overline{M}$ and $\overline{N} \subseteq \overline{N}^{(r)}$.

The cone σ considered with respect to the new lattice $\overline{M}_{(r)}$ is a Gorenstein cone of index 1, and the reflexive polytope $\sigma_{(r)} = r\tilde{\Delta}$ is its support. Thus, we obtain:

Proposition 1.15. *Let $\tilde{\Delta} \subseteq \widetilde{M}_{\mathbb{R}}$ be a Gorenstein polytope of index r and $\tilde{\Delta}^* \subset \widetilde{N}_{\mathbb{R}}$ be the dual Gorenstein polytope. If $\sigma \subseteq \overline{M}_{\mathbb{R}}$ is a reflexive Gorenstein cone of index r with support $\tilde{\Delta} \subseteq \widetilde{M}_{\mathbb{R}}$, then the dual reflexive polytope $(r\tilde{\Delta})^*$ equals $\tilde{\Delta}^*$ with respect to the refined affine lattice*

$$\overline{N}^{(r)} \cap \{y \in N_{\mathbb{R}} : \langle m_{\sigma^{\vee}}, y \rangle = 1\}.$$

Proposition 1.16. *Let ρ be a facet of $\sigma^{\vee}$. We denote by $\text{lin}(\rho)$ the minimal linear subspace in $\overline{N}_{\mathbb{R}}$ containing ρ . Then*

$$\text{lin}(\rho) \cap \overline{N} = \text{lin}(\rho) \cap \overline{N}^{(r)}.$$

In particular, one has

$$\partial\sigma^{\vee} \cap \overline{N} = \partial\sigma^{\vee} \cap \overline{N}^{(r)}$$

and

$$\partial\tilde{\Delta}^* \cap \tilde{N} = \partial(r\tilde{\Delta})^* \cap \overline{N}^{(r)}.$$

Proof. Let $x \in \text{lin}(\rho) \cap \overline{N}^{(r)}$. Since ρ is a facet of $\sigma^{\vee}$, there is a vertex v of $\tilde{\Delta}$ such that $\langle v, y \rangle = 0$ for all $y \in \rho$. From $\langle v, n_{\sigma} \rangle = 1$ we get $\overline{N} = \mathbb{Z}n_{\sigma} \oplus (\overline{N} \cap v^{\perp})$. Hence $\overline{N}^{(r)} = \frac{1}{r}\mathbb{Z}n_{\sigma} \oplus (\overline{N} \cap v^{\perp})$. Now, $\langle v, x \rangle = 0$ implies $x \in \overline{N}$. $\square$

Remark 1.17. From the combinatorial point of view, the duality of Gorenstein polytopes “coincides” with the polar duality. However, there exists a “subtlety” coming from the occurring lattices. Let $\tilde{\Delta}$ be a Gorenstein polytope of index $r > 1$. Then the Gorenstein polytope $\tilde{\Delta}^*$ has no interior lattice points. Proposition 1.16 shows that the reflexive polytope $(r\tilde{\Delta})^*$ and $\tilde{\Delta}^*$ have the same sets of boundary lattice points. Assume that r is not a prime number. Let $k > 1$ be any proper divisor of r . Then $k\tilde{\Delta}$ is a Gorenstein polytope of index r/k . The dual Gorenstein polytope $(k\tilde{\Delta})^*$ has the same lattice points on the boundary as $(r\tilde{\Delta})^*$ and $\tilde{\Delta}^*$. Therefore,

two Gorenstein polytopes $(k\tilde{\Delta})^*$ and $\tilde{\Delta}^*$ have the same combinatorial structure and also the same lattice point structure. However, $(k\tilde{\Delta})^*$ and $\tilde{\Delta}^*$ are Gorenstein polytopes of different index, so they are *not* isomorphic as lattice polytopes.

Consider an example of this phenomenon:

Example 1.18. Let Δ be a unimodular 3-dimensional simplex, i.e., it has lattice volume 1. Then Δ is a Gorenstein polytope of index 4. We have $\Delta \cong \Delta^*$, i.e. Δ is a selfdual Gorenstein polytope. The dual reflexive polytope $(4\Delta)^*$ is a 3-dimensional reflexive simplex with 4 lattice points on the boundary. Now consider 2Δ as a Gorenstein polytope of index 2. The dual Gorenstein polytope $(2\Delta)^*$ is a Gorenstein simplex of index 2 having lattice volume 2 and having only 4 vertices as lattice points (the same number of lattice points as in Δ^*).

The polytope $4(\Delta^*)$ is a reflexive simplex with 34 lattice points on the boundary. The polytope $2(2\Delta)^*$ is a self-dual reflexive simplex with 10 lattice points on the boundary. Pictures of these polytopes are given in Example 2.17 where the polytope 2Δ is denoted by $\tilde{\Delta}^*$.

2. CAYLEY POLYTOPES AND SPECIAL SIMPLICES

In this section we consider Cayley polytopes of length r and give a criterion for a Gorenstein polytope $\tilde{\Delta}$ of index r to be a Cayley polytope of length r . For this purpose we use the notion of a special $(r-1)$ -dimensional simplex in the dual Gorenstein polytope $\tilde{\Delta}^*$.

2.1. Cayley polytopes. Let M, N be dual lattices of rank d ; and let $\Delta_1, \dots, \Delta_r \subseteq M_{\mathbb{R}}$ be lattice polytopes.

Definition 2.1. Consider the lattice $\overline{M} := M \oplus \mathbb{Z}^r$, where $e_1, \dots, e_r$ is the standard lattice basis of $\mathbb{Z}^r$. Then $\overline{M}$ contains the affine sublattice $\widetilde{M} := M \oplus (\text{aff}(e_1, \dots, e_r) \cap \mathbb{Z}^r)$. We identify $\Delta_i \subseteq M_{\mathbb{R}}$ with $\tilde{\Delta}_i := \Delta_i \times e_i \subseteq \widetilde{M}_{\mathbb{R}}$ for $i = 1, \dots, r$. The polytope $\Delta_1 * \dots * \Delta_r := \text{Conv}(\tilde{\Delta}_1, \dots, \tilde{\Delta}_r) \subseteq \widetilde{M}_{\mathbb{R}}$ is called the *Cayley polytope of length r* associated to $\Delta_1, \dots, \Delta_r$. The cone

$$\mathbb{R}_{\geq 0}(\Delta_1 * \dots * \Delta_r) = \mathbb{R}_{\geq 0}\tilde{\Delta}_1 + \dots + \mathbb{R}_{\geq 0}\tilde{\Delta}_r \subseteq \overline{M}$$

is called the associated *Cayley cone*.

If $\text{aff}(\Delta_1, \dots, \Delta_r) = M_{\mathbb{R}}$, then the associated Cayley cone is a Gorenstein cone of dimension $d+r$, and $\tilde{\Delta} = \Delta_1 * \dots * \Delta_r$ is its support.

Definition 2.2. Let $e_1, \dots, e_r \in \mathbb{Z}^r$ be the standard lattice basis. Then $S_{r-1} := \text{Conv}(e_1, \dots, e_r)$ is called an *unimodular simplex* of dimension $r-1$.

Proposition 2.3. Let $\sigma \subseteq \overline{M}_{\mathbb{R}}$ be a Gorenstein cone with support $\tilde{\Delta} \subseteq \widetilde{M}_{\mathbb{R}}$. Then the following statements are equivalent:

- (1) σ is a Cayley cone associated to r lattice polytopes;
- (2) $\tilde{\Delta}$ is a Cayley polytope of length r ;

(3) *There is a lattice projection $\widetilde{M} \rightarrow \mathbb{Z}^r$, which maps $\tilde{\Delta}$ surjectively on S_{r-1} ;*

(4) *There are nonzero $e_1^*, \dots, e_r^* \in \sigma^\vee \cap \overline{N}$, such that $e_1^* + \dots + e_r^* = n_\sigma$.*

Moreover, the lattice vectors $e_1^, \dots, e_r^*$ in (4) form a part of a basis of $\overline{N}$ and the Cayley structure of $\tilde{\Delta}$ is uniquely determined by r polytopes*

$$\tilde{\Delta}_i := \{x \in \tilde{\Delta} : \langle x, e_j^* \rangle = 0 \text{ for } j \neq i\} \quad (i = 1, \dots, r).$$

These polytopes have properties $\langle \tilde{\Delta}_i, e_i^ \rangle = 1$, and $\tilde{\Delta} = \text{Conv}(\tilde{\Delta}_1, \dots, \tilde{\Delta}_r)$.*

Proof. (1), (2), (3) are obviously equivalent.

(1) $\Rightarrow$ (4): Let $b_1, \dots, b_d, e_1, \dots, e_r$ be a basis of the lattice $\overline{M}$, where $b_1, \dots, b_d$ is a basis of the lattice M and $e_1, \dots, e_r$ the standard basis of $\mathbb{Z}^r$. Consider the dual basis $b_1^*, \dots, b_d^*, e_1^*, \dots, e_r^*$ of the lattice $\overline{N}$. Then we have $\langle x_j, e_i^* \rangle = \delta_{ij}$ for $i, j \in \{1, \dots, r\}$ and $x_j \in \tilde{\Delta}_j$ (δ_{ij} denotes the Kronecker symbol, i.e., $\delta_{ij} = 1$, for $i = j$, and 0 otherwise). Therefore we obtain $e_1^*, \dots, e_r^* \in \sigma^\vee$, $\langle \tilde{\Delta}, e_1^* + \dots + e_r^* \rangle = 1$. The latter implies $e_1^* + \dots + e_r^* = n_\sigma$.

(4) $\Rightarrow$ (1): Define the polytopes $\tilde{\Delta}_i := \{x \in \tilde{\Delta} : \langle x, e_j^* \rangle = 0 \text{ for } j \neq i\}$ for $i = 1, \dots, r$. Let x be a vertex of $\tilde{\Delta}$. Then $1 = \langle x, e_1^* + \dots + e_r^* \rangle$. Since $0 \leq \langle x, e_i^* \rangle \in \mathbb{Z}$ for $i = 1, \dots, r$, we get that $x \in \tilde{\Delta}_i$ for some i and $\langle x, e_i^* \rangle = 1$. Hence $\tilde{\Delta} = \text{Conv}(\tilde{\Delta}_1, \dots, \tilde{\Delta}_r)$. Take any $i \in \{1, \dots, r\}$. We claim that $\tilde{\Delta}_i \neq \emptyset$. Indeed, otherwise we would have $\langle \tilde{\Delta}, e_i^* \rangle = 0$. This would contradict the condition $e_i^* \neq 0$. Hence the mapping $x \rightarrow (\langle x, e_1^* \rangle, \dots, \langle x, e_r^* \rangle)$ gives a projection of lattices $\widetilde{M} \rightarrow \mathbb{Z}^r$ such that the image of $\tilde{\Delta}$ is S_{r-1} . In particular, $e_1^*, \dots, e_r^*$ form a part of a basis of $\overline{N}$. $\square$

Recall the following definition from [BB97, Def. 3.9]:

Definition 2.4. A reflexive Gorenstein cone σ of index r is said to be *completely split*, if σ is the Cayley cone associated to r lattice polytopes.

By 2.3, we obtain the following characterization of completely split reflexive Gorenstein cones:

Corollary 2.5. *Let $\sigma \subseteq \overline{M}_{\mathbb{R}}$ be a reflexive Gorenstein cone of index r with support $\tilde{\Delta}$. Then the following statements are equivalent:*

- (1) *σ is completely split;*
- (2) *there exist lattice points $e_1^*, \dots, e_r^* \in \tilde{\Delta}^* \cap \overline{N}$ such that*

$$e_1^* + \dots + e_r^* = n_\sigma.$$

Moreover, there is a 1-to-1 correspondence between all possible collections $\{e_1^, \dots, e_r^*\}$ of r lattice points in $\tilde{\Delta}^*$ such that $e_1^* + \dots + e_r^* = n_\sigma$ and all possible Cayley polytope structures of length r of the Gorenstein polytope $\tilde{\Delta}$.*

Proof. (1) $\Rightarrow$ (2): Assume that σ is completely split. By 2.3(4) there exist lattice points $e_1^*, \dots, e_r^* \in \sigma^\vee$ such that $e_1^* + \dots + e_r^* = n_\sigma$. Since $r =$

$\langle m_{\sigma^\vee}, n_\sigma \rangle = \langle m_{\sigma^\vee}, e_1^* \rangle + \cdots + \langle m_{\sigma^\vee}, e_r^* \rangle$ and $1 \leq \langle m_{\sigma^\vee}, e_i^* \rangle \in \mathbb{Z}$ for all $i = 1, \dots, r$, we get $\langle m_{\sigma^\vee}, e_i^* \rangle = 1$ for all $i = 1, \dots, r$. Thus, all lattice points $e_1^*, \dots, e_r^*$ belong to $\tilde{\Delta}^*$.

(2) $\Rightarrow$ (1): The statement follows immediately from 2.3, because the index of σ equals $\langle m_{\sigma^\vee}, n_\sigma \rangle = \langle m_{\sigma^\vee}, e_1^* \rangle + \cdots + \langle m_{\sigma^\vee}, e_r^* \rangle = r$. $\square$

By 2.3, every completely split reflexive Gorenstein cone $\sigma \subset \overline{M}_{\mathbb{R}}$ determines r lattice polytopes $\tilde{\Delta}_1, \dots, \tilde{\Delta}_r$. On the other hand, one can start with r lattice polytopes $\Delta_1, \dots, \Delta_r \subset M_{\mathbb{R}}$ and ask whether the corresponding Cayley cone is reflexive of index r , or whether the Cayley polytope $\Delta_1 * \cdots * \Delta_r$ is a Gorenstein polytope of index r . The answer is contained in [BB97, Prop.3.6]:

Theorem 2.6. *Let $\Delta_1, \dots, \Delta_r \subset M_{\mathbb{R}}$ be lattice polytopes such that the Minkowski sum $\Delta_1 + \cdots + \Delta_r$ has dimension d . Then the following statements are equivalent:*

- (1) *The associated Cayley cone σ is reflexive of index r ;*
- (2) *$\Delta_1 * \cdots * \Delta_r$ is a Gorenstein polytope of index r ;*
- (3) *$\Delta_1 + \cdots + \Delta_r$ is a reflexive polytope (with interior lattice point m).*

In this case, the lattice point $m_{\sigma^\vee} \in \sigma$ (see Def. 1.10) equals $m \times (e_1 + \cdots + e_r)$.

For convenience, we give a purely combinatorial proof of this statement.

Proof. (1) $\Leftrightarrow$ (2): Follows from Prop. 1.11.

(2) $\Rightarrow$ (3): Let $\tilde{\Delta} = \Delta_1 * \cdots * \Delta_r$ be a Gorenstein polytope of index r . Denote by π the projections $\overline{M} \rightarrow \mathbb{Z}^r$ and $\overline{M}_{\mathbb{R}} \rightarrow \mathbb{R}^r$. Since $r\tilde{\Delta}$ is reflexive, there exists a unique lattice point $m \in \text{int}(r\tilde{\Delta})$. Moreover, the π -image of m can only be the unique interior point $s_1 := (1, \dots, 1)$ in the reflexive simplex rS_{r-1} (where $S_r = \text{Conv}(e_1, \dots, e_r)$). We remark that the intersection of the affine subspace $\pi^{-1}(s_1)$ with $r\tilde{\Delta}$ is exactly $\Delta_1 + \cdots + \Delta_r$. Since m has integral distance 1 from any facet of $r\tilde{\Delta}$, the lattice point m has also integral distance 1 from any facet of $\Delta_1 + \cdots + \Delta_r$ (facets of $\Delta_1 + \cdots + \Delta_r$ are intersections of facets of $r\tilde{\Delta}$ with the affine subspace $\pi^{-1}(s_1)$).

(3) $\Rightarrow$ (1): Assume that the lattice polytope $\Delta := \Delta_1 + \cdots + \Delta_r$ is reflexive. Denote by m the unique interior lattice point in P . Consider the Cayley cone σ over $\tilde{\Delta} = \Delta_1 * \cdots * \Delta_r$ and denote by $m_{\sigma^\vee}$ the lattice point $m \times (e_1 + \cdots + e_r)$ in $\text{int}(\sigma)$. We have to show that $\langle m_{\sigma^\vee}, z \rangle = 1$ for any primitive lattice vector $z \in \overline{N}$ generating a 1-dimensional face $\mathbb{R}_{\geq 0}z$ of the dual cone $\sigma^\vee \subset \overline{N}_{\mathbb{R}}$.

Let $\Gamma := z^\perp \cap \sigma$ the facet of σ which is orthogonal to z . There are the following two possibilities:

Case 1. There exists $i \in \{1, \dots, r\}$ such that $\Gamma \cap (\Delta_i \times e_i) = \emptyset$. Then Γ is orthogonal to e_i^* , because Γ is generated by some lattice points from the polytopes $(\Delta_j \times e_j)$ ($j = 1, \dots, r$) and $\langle x, e_i^* \rangle = \delta_{ij}$ for $x \in (\Delta_j \times e_j)$. Since e_i^* is primitive and $e_i^* \in \sigma^\vee$, we get $z = e_i^*$ and $\langle m_{\sigma^\vee}, z \rangle = \langle e_1 + \cdots + e_r, e_i^* \rangle = 1$.

Case 2. For any $i \in \{1, \dots, r\}$ there exists a lattice point $p_i \in \Delta_i \times e_i$ which is contained in Γ . In this case, Γ is a Cayley cone of r lattice polytopes $P_i := \Gamma \cap (\Delta_i \times e_i)$ ($i = 1, \dots, r$) such that $p_i \in P_i$ ($1 \leq i \leq r$). Moreover, $P_\Gamma := \Gamma \cap (\Delta \times (e_1 + \dots + e_r)) = P_1 + \dots + P_r$ is a $(d-1)$ -dimensional face of $\Delta \times (e_1 + \dots + e_r) \cong \Delta_1 + \dots + \Delta_r$.

Let M' be the $(d+1)$ -dimensional sublattice in $\overline{M}$ defined by $r-1$ equations $\langle x, e_1^* \rangle = \dots = \langle x, e_r^* \rangle$. Then $\sigma_\Delta := M_{\mathbb{R}}' \cap \sigma$ is exactly the cone over the d -dimensional reflexive polytope $\Delta = \Delta_1 + \dots + \Delta_r$. Moreover, $m_{\sigma^\vee}$ is the corresponding interior lattice point in $\text{int}(\sigma_\Delta)$. There exists a primitive lattice vector z' in the dual lattice N' such that $z' \in \sigma_\Delta^\vee$, $(z')^\perp \cap \sigma_\Delta = \Gamma \cap \sigma_\Delta$, and $\langle m_{\sigma^\vee}, z' \rangle = 1$. We identify the dual lattice N' with $\overline{N} / \sum_{i,j} \mathbb{Z}(e_i^* - e_j^*)$. Since z and z' have the same orthogonal subspace in $\overline{M}'$ there exists a positive integer k such that $z = kz'$ modulo the sublattice generated by $e_2^* - e_1^*, \dots, e_r^* - e_1^*$. In this case, we have $\langle m_{\sigma^\vee}, z \rangle = k \langle m_{\sigma^\vee}, z' \rangle = k$, because $\langle m_{\sigma^\vee}, e_i^* - e_j^* \rangle = 1 - 1 = 0$ for all i, j . Now it remains to show that $k = 1$. Using the splittings

$$\overline{N} = N \oplus \bigoplus_{i=1}^r \mathbb{Z}e_i^*, \quad \bigoplus_{i=1}^r \mathbb{Z}e_i^* = \mathbb{Z}e_1^* \oplus \bigoplus_{i=2}^r \mathbb{Z}(e_i^* - e_1^*),$$

we can write

$$z = kz' + \sum_{i=1}^r l_i e_i^*$$

for some integers $l_1, \dots, l_r$. Applying $\langle p_j, * \rangle$ to both sides of the last equation, we get

$$0 = \langle p_j, z \rangle = \langle p_j, kz' \rangle + \langle p_j, \sum_{i=1}^r l_i e_i^* \rangle = k \langle p_j, z' \rangle + l_j,$$

i.e., k divides l_j for all $1 \leq j \leq r$. Since $z = kz' + \sum_{i=1}^r l_i e_i^*$ is a primitive lattice vector, we get $k = 1$. $\square$

2.2. Special simplices. Our next purpose is to explain the relations between Cayley cones and Cayley polytopes to some results in [Ath05, BR05]. It would be more convenient to work with polytopes in the dual space $N_{\mathbb{R}}$. For this reason, we denote these polytopes by ∇ .

Athanasiadis introduced in [Ath05] the concept of a special simplex S of a convex polytope ∇ . Here we slightly modify this definition by demanding the vertices of the simplex S to be lattice points in ∇ .

Definition 2.7. Let $\nabla \subset N_{\mathbb{R}}$ be a d -dimensional lattice polytope. A simplex S spanned by r affinely independent lattice points in ∇ is called a *special* $(r-1)$ -simplex of ∇ , if each facet of ∇ contains exactly $r-1$ vertices of S .

Now, we can give an addendum to Cor. 2.5:

Proposition 2.8. *Let $\nabla \subset N_{\mathbb{R}}$ be a Gorenstein polytope of index r . There is a 1-to-1 correspondence between Cayley polytope structures of length r of the dual Gorenstein polytope $\nabla^* \subset M_{\mathbb{R}}$ and special $(r-1)$ -simplices $S \subset \nabla$. A special $(r-1)$ -simplex $S \subset \nabla$ can be characterized as a lattice $(r-1)$ -dimensional simplex in ∇ , which is not contained in the boundary of ∇ . Moreover, all special $(r-1)$ -simplices in ∇ have the same barycenter.*

Proof. Let $\sigma \subseteq M_{\mathbb{R}}$ be the reflexive Gorenstein cone associated to $\Delta := \nabla^*$. We identify Δ with the support $\tilde{\Delta}$ of σ and ∇ with the support $\tilde{\nabla}$ of $\sigma^\vee$. If Δ is a Cayley polytope of length r , then, by Cor. 2.5, there exist $n_1, \dots, n_r \in \tilde{\nabla} \cap \overline{N}$ such that $n_1 + \dots + n_r = n_\sigma$. If F is a facet of $\tilde{\nabla}$, then there is a vertex $v \in \tilde{\Delta}$ such that $\langle v, F \rangle = 0$. Let $\tilde{\Delta} = \text{Conv}(\tilde{\Delta}_1, \dots, \tilde{\Delta}_r)$ as in Prop. 2.3. Then $v \in \mathcal{V}(\tilde{\Delta}_i)$ for some i . This implies $\langle v, n_i \rangle = 1$ and $\langle v, n_j \rangle = 0$ for $j \neq i$. So $n_i \notin F$, but $n_j \in F$ for $j \neq i$. So $\text{Conv}(n_1, \dots, n_r)$ is a special $(r-1)$ -simplex in ∇ .

Now assume that $n_1, \dots, n_r \in \tilde{\nabla} \cap \overline{N}$ are vertices of a $(r-1)$ -dimensional simplex that is not contained in the boundary of ∇ . Then $n_1 + \dots + n_r \in r\tilde{\nabla}$ is in the interior of $\sigma^\vee$. Since $r\tilde{\nabla}$ is reflexive, we have $\text{int}(r\tilde{\nabla}) \cap \overline{N} = \{n_\sigma\}$. Therefore $n_1 + \dots + n_r = n_\sigma$. By Prop. 2.3, Δ is a Cayley polytope of length r . Moreover, above arguments show that $S = \text{Conv}(n_1, \dots, n_r)$ is a special $(r-1)$ -simplex in ∇ . If $S' = \text{Conv}(n'_1, \dots, n'_r)$ is another special simplex in ∇ then

$$n_1 + \dots + n_r = n'_1 + \dots + n'_r = n_\sigma.$$

Thus S and S' have the same barycenter. $\square$

Example 2.9. If $\nabla \subset N_{\mathbb{R}}$ is a reflexive polytope (Gorenstein polytope of index $r = 1$) then there exists only one special $(r-1)$ -simplex in ∇ which is exactly the unique interior lattice point in ∇ .

Example 2.10. Let ∇_d be the standard d -dimensional unit cube in $\mathbb{R}^d$: $\nabla_d := \{x \in \mathbb{R}^d : 0 \leq x_i \leq 1\}$. Then ∇_d is a Gorenstein polytope of index $r = 2$. There exist exactly 2^{d-1} different special 1-simplices in ∇_d which are spanned by all possible pairs of opposite vertices of ∇_d . The dual Gorenstein polytope ∇_d^* has exactly d special 1-simplices which 1-to-1 correspond to d different Cayley structures of the d -dimensional unit cube ∇_d .

Let us recall the following definition:

Definition 2.11. A lattice polytope $\nabla \subseteq N_{\mathbb{R}}$ is called *integrally closed*, if any lattice point in the Gorenstein cone σ over ∇ is a sum of lattice points from the support $\sigma_{(1)} \cong \nabla$.

We get the following corollary:

Corollary 2.12. *Let $\nabla \subseteq N_{\mathbb{R}}$ be an integrally closed Gorenstein polytope of index r . Then ∇ contains a special $(r-1)$ -simplex and ∇^* is a Cayley polytope of length r .*

Proof. It follows immediately from the fact that the lattice point n_σ is a sum of r lattice points $n_1, \dots, n_r$ from $\tilde{\nabla}$. $\square$

Remark 2.13. One can easily show that if a lattice polytope ∇ admits a unimodular triangulation, then ∇ is integrally closed. Under the assumption that ∇ admits a unimodular triangulation the statement in 2.12 was proved in [Sti98, Prop. 3].

One of the main results in [BR05] is Corollary 7, where Bruns and Roemer show that for any integrally closed Gorenstein polytope there exists a reflexive polytope with the same h^* -vector. Here we give a slightly more general version of this result that does not need any commutative algebra. We first recall the definition of the h^* -polynomial (also called δ -polynomial), cf. [BN06, Hib92, Sta87]:

Definition 2.14. Let $\nabla \subset N_{\mathbb{R}}$ be a d -dimensional lattice polytope. The h^* -polynomial $h_{\nabla}^*(t)$ of ∇ is defined as the numerator of the Ehrhart series, i.e., $h_{\nabla}^*(t) = (1-t)^{d+1} \sum_{k \in \mathbb{N}} |k\nabla \cap N| t^k$.

Remark 2.15. The h^* -polynomial has degree at most d . It is well-known, see [Hib92], that ∇ is Gorenstein of index r if and only if for the h^* -polynomial one has:

$$t^{d-r+1} h_{\nabla}^*(t^{-1}) = h_{\nabla}^*(t).$$

In this case, ∇ is reflexive if and only if the degree of h_{∇}^* is d .

The following theorem is very close to results in [Ath05], [BR05], and [OH05]:

Theorem 2.16. Let $\nabla \subseteq N_{\mathbb{R}}$ be a Gorenstein polytope of index r . Assume that ∇ contains a special $(r-1)$ -simplex with vertices $n_1, \dots, n_r$, i.e. the dual Gorenstein polytope ∇^* is a Cayley polytope associated with some lattice polytopes $\Delta_1, \dots, \Delta_r$. Then the projection of ∇ along $\text{aff}(n_1, \dots, n_r)$ is a $(d-r+1)$ -dimensional reflexive polytope P such that $P = (\Delta_1 + \dots + \Delta_r)^*$ and

$$h_P^*(t) = h_{\nabla}^*(t).$$

Proof. The statement follows straightforward from [BR05, Lemma 4, Lemma 5]. Another variant of the same ideas is contained in [RW05, Prop.3.12] and [Ath05, Lemma 3.4, Cor. 4.1, Cor. 4.2]. For convenience, we explain the main combinatorial idea of the proof.

Let $\tilde{\nabla}$ be the support of the Gorenstein cone over ∇ . We identify ∇ with $\tilde{\nabla}$. For any $i \in \{1, \dots, r\}$ let $\mathcal{F}_i$ be the set of facets of $\tilde{\nabla}$ that do *not* contain n_i . We denote by Γ the union of all faces $F_1 \cap \dots \cap F_r \subset \tilde{\nabla}$, where $F_i \in \mathcal{F}_i$ for $i = 1, \dots, r$.

Take one such a face $G := F_1 \cap \dots \cap F_r \in \Gamma$. We define a lattice polytope $G' := \text{Conv}(G, n_1, \dots, n_r)$. Let $v_1, \dots, v_r \in \tilde{\nabla}^*$ be the vertices corresponding

to the facets $F_1, \dots, F_r$. By Prop. 2.3, we have $\langle v_j, n_i \rangle = \delta_{i,j}$ for $i, j \in \{1, \dots, r\}$. Therefore G' is an r -fold pyramid over G , i.e., $h_{G'}^*(t) = h_G^*(t)$, cf. [Bat07, Remark 2.6]. By [BR05, Lemma 4] the set of all G' (together with their subfaces) form a polyhedral decomposition of $\tilde{\nabla}$.


Denote by π the projection of ∇ along $\text{aff}(n_1, \dots, n_r)$. Let $x \in \text{int}(\nabla)$ be the unique interior point with lattice distance $1/r$ from any facet of ∇ . We define the refined lattice $N' \supseteq N$ with $r = [N' : N]$ by $N' := N + \mathbb{Z}x$. By Prop. 2.8 and Prop. 2.3, we have $(n_1 + \dots + n_r)/r = x$. So the projection π can be seen as the surjective map $N' \rightarrow N'/(\mathbb{Z}x + \sum_{i=1}^{r-1} \mathbb{Z}n_i)$ where $x, n_1, \dots, n_{r-1}$ form a part of $\mathbb{Z}$ -basis of N' . Let F be a face of $\tilde{\nabla}$ which is the preimage of a facet of P . By definition of special $(r-1)$ -simplex, any facet of $\tilde{\nabla}$ is contained in $\mathcal{F}_1 \cup \dots \cup \mathcal{F}_r$. Since $n_i \notin F$ for all $i = 1, \dots, r$ ($\pi(n_i) = 0 \notin \partial P$), there exists $F_i \in \mathcal{F}_i$ which contains F . Thus $F \subseteq G = F_1 \cap \dots \cap F_r$ for some $G \in \Gamma$. On the other hand, $\dim F \geq d-r$ and $\dim G = d-r$. Therefore, $F = G$. By [BR05, Lemma 5], the projection $\pi : \tilde{\nabla} \rightarrow P$ respects two polyhedra decompositions

$$\tilde{\nabla} = \bigcup_{G \in \Gamma} \text{Conv}(G, n_1, \dots, n_r)$$

$$P = \bigcup_{G \in \Gamma} \text{Conv}(\pi(G), 0),$$

where each $\text{Conv}(G, n_1, \dots, n_r)$ is a r -fold pyramid over a face $G \in \Gamma$ and the polytope $\text{Conv}(\pi(G), 0)$ is 1-fold pyramid with vertex 0 over a facet in $\pi(G) \subset \partial P$. Since the h^* -polynomials of $\text{Conv}(G, n_1, \dots, n_r)$ and $\text{Conv}(\pi(G), 0)$ are the same, a standard inclusion-exclusion argument shows that the h^* -polynomials of $\tilde{\nabla}$ and P are the same. Using 2.15, this implies that P is a reflexive polytope. By Theorem 2.6, $\Delta_1 + \dots + \Delta_r$ is a reflexive polytope. Moreover, the proof of Theorem 2.6 shows that P is the dual reflexive polytope to $\Delta_1 + \dots + \Delta_r$. $\square$

Example 2.17. The above theorem is illustrated in the following example: Consider one-dimensional lattice polytopes $\Delta_1 := \text{Conv}((-1, 0), (0, -1))$ and $\Delta_2 := \text{Conv}((0, 0), (1, 1))$. We remark that Δ_2 contains 0, but Δ_1 does not. The Cayley polytope $\tilde{\Delta} = \Delta_1 * \Delta_2$ is a Gorenstein polytope of index 2. On the other hand, $\tilde{\Delta}^*$ is isomorphic to $2S_3$, where S_3 is the unimodular 3-simplex. The polytope $\tilde{\Delta}^*$ contains a special 1-simplex S corresponding to the Cayley polytope structure of $\tilde{\Delta}$ (the two vertices of S are marked by boxes). By projecting along this simplex, we obtain the reflexive square P . The boundary subcomplex Γ of $\tilde{\Delta}^*$ (which maps bijectively to the boundary of P) is marked by thick edges. The dual reflexive polytope P^* of P is $\Delta_1 + \Delta_2$. It is the intersection of $2\tilde{\Delta}$ with an affine plane which is orthogonal to the special simplex S .



3. COMBINATORICS OF NEF-PARTITIONS

3.1. Characterization of nef-partitions. Now we are interested in the case when both dual reflexive Gorenstein cones are completely split. Using special simplices, we want to obtain a simple combinatorial characterization of nef-partitions and their duality.

Let us consider the following purely combinatorial definition of a nef-partition (cf. [KRS03, Prop.3.2]):

Definition 3.1. Let $\Delta \subset M_{\mathbb{R}}$ be a d -dimensional reflexive polytope and $m = \text{int}(\Delta) \cap M$. A Minkowski sum decomposition $\Delta = \Delta_1 + \cdots + \Delta_r$ where $\Delta_1, \dots, \Delta_r \subseteq M_{\mathbb{R}}$ are lattice polytopes is called a *nef-partition* of Δ of length r , if there exist lattice points $p_1 \in \Delta_1, \dots, p_r \in \Delta_r$ such that $p_1 + \cdots + p_r = m$. Since the reflexive polytope Δ is determined by $\Delta_1, \dots, \Delta_r$, for simplicity, we use the name *nef-partition* just for the set of lattice polytopes $\Delta_1, \dots, \Delta_r$.

We call a nef-partition *centered*, if $p_i = 0$ for all $1 \leq i \leq r$. We call a nef-partition *proper*, if $\dim \Delta_i > 0$ for all $1 \leq i \leq r$.

Example 3.2. Let $\Delta \subset \mathbb{R}^d$ be the d -dimensional reflexive cube defined by the conditions $|x_i| \leq 1$ ($1 \leq i \leq d$). Then Δ is a Minkowski sum of two unit cubes Δ_1 and Δ_2 defined as

$$\Delta_1 := \{x = (x_i) \in \mathbb{R}^d : 0 \leq x_i \leq 1 \forall i\},$$

$$\Delta_2 := \{x = (x_i) \in \mathbb{R}^d : -1 \leq x_i \leq 0 \forall i\}.$$

Since $0 \in \Delta_i$ ($i = 1, 2$), the Minkowski sum $\Delta = \Delta_1 + \Delta_2$ defines a proper centered nef-partition of length 2. This example comes from the Gorenstein polytopes considered in 2.10.

Remark 3.3. It clear that if $\Delta = \Delta_1 + \cdots + \Delta_r$ is a nef-partition of Δ , then

$$(\Delta - m) = (\Delta_1 - p_1) + \cdots + (\Delta_r - p_r)$$

is a centered nef-partition. Moreover, if

$$\Delta = \sum_{i=1}^r \Delta_i$$

is a centered nef-partition of Δ , then

$$\Delta = \sum_{i: \Delta_i \neq 0} \Delta_i$$

is a proper centered nef-partition. Therefore, the study of arbitrary nef-partitions immediately reduces to study of proper centered nef-partitions.

Remark 3.4. The above definition of proper centered nef-partitions is nearly precisely the one given by Kreuzer, Riegler and Sahakyan in [KRS03, Prop.3.2]. They also demand an additional assumption $\Delta_i \cap \Delta_j = \{0\}$ for $i \neq j$, which is actually unnecessary (see Cor. 3.17).

Remark 3.5. From the viewpoint of algebraic geometry, proper centered nef-partitions were introduced by Borisov in [Bor93]. Let $\mathbb{P}_\Delta$ be a Gorenstein toric Fano variety corresponding to the reflexive polytope Δ which contains 0 in its interior. Denote by $\mathcal{T}$ the d -dimensional torus acting on $\mathbb{P}_\Delta$ (we identify $\mathcal{T}$ with its open dense orbit in $\mathbb{P}_\Delta$). A Minkowski sum decomposition $\Delta = \Delta_1 + \cdots + \Delta_r$ describes the anticanonical sheaf $\mathcal{O}(-K_{\mathbb{P}_\Delta})$ as a tensor product of r $\mathcal{T}$ -equivariant semi-ample line bundles $\mathcal{L}_1, \dots, \mathcal{L}_r$. In this case, the lattice points $p_i \in \Delta_i$ can be identified with some global $\mathcal{T}$ -invariant sections s_i of $\mathcal{L}_i$ which do not vanish on $\mathcal{T} \subset \mathbb{P}_\Delta$. The condition $p_1 + \cdots + p_r = 0$ means that if D_i is the zero set of the global section s_i ($i = 1, \dots, r$), then $D_1 + \cdots + D_r$ is the anticanonical divisor of $\mathbb{P}_\Delta$ which contains every irreducible component of $\mathbb{P}_\Delta \setminus \mathcal{T}$ with multiplicity 1. In particular, D_i and D_j do not have common irreducible components for $i \neq j$. Moreover, $\dim \Delta_i \neq 0$ if and only if $D_i \neq 0$. Therefore, if $\dim \Delta_i \neq 0$ for all $1 \leq i \leq r$, then we get a *partition* of the set of irreducible components of $\mathbb{P}_\Delta \setminus \mathcal{T}$ into r nonempty subsets which define *nef-divisors* $D_1, \dots, D_r$ such that $D_1 + \cdots + D_r$ is the anticanonical divisor. This explains the origin of the notion *nef-partition* in [Bor93].

We characterize nef-partitions in the following way:

Proposition 3.6. *Let Δ be a reflexive polytope with the unique interior lattice point m , and let $\Delta_1, \dots, \Delta_r$ be lattice polytopes such that $\Delta = \Delta_1 + \cdots + \Delta_r$. Denote by σ the Cayley cone associated to $\Delta_1, \dots, \Delta_r$. Then the following statements are equivalent:*

- (1) *the dual cone $\sigma^\vee$ is a completely split reflexive Gorenstein cone of index r*

- (2) $\Delta_1 * \cdots * \Delta_r$ is a Gorenstein polytope of index r containing a special $(r-1)$ -simplex
- (3) there exist lattice points $p_i \in \Delta_i$ ($1 \leq i \leq r$) such that $p_1 + \cdots + p_r = m$, i.e., $\Delta = \Delta_1 + \cdots + \Delta_r$ is a nef-partition.

Proof. (1) $\Rightarrow$ (2): If $\sigma^\vee$ is a reflexive Gorenstein cone of index r , then this holds also for σ . Hence Theorem 2.6 implies that $\Delta_1 * \cdots * \Delta_r$ is a Gorenstein polytope of index r . Since $\sigma^\vee$ is completely split, Prop. 2.8 implies that $\Delta_1 * \cdots * \Delta_r$ contains a special $(r-1)$ -simplex.

(2) $\Rightarrow$ (3): Since $\sigma^\vee$ is completely split, Cor. 2.5 implies that there are $\tilde{p}_1, \dots, \tilde{p}_r \in \tilde{\Delta} \cap \overline{M}$ such that $\tilde{p}_1 + \cdots + \tilde{p}_r = m_{\sigma^\vee}$. Let $e_1^*, \dots, e_r^* \in \tilde{\Delta}^* \cap \overline{N}$ as in Prop. 2.3 and Cor. 2.5. Since $\langle m_{\sigma^\vee}, e_i^* \rangle = 1 \ \forall i$ and $\langle \tilde{p}_j, e_i^* \rangle$ is a nonnegative integer $\forall j$, every polytope $\tilde{\Delta}_i = \Delta_i \times e_i$ contains exactly one point from $\{\tilde{p}_1, \dots, \tilde{p}_r\}$. Without loss of generality we can assume that $\tilde{p}_i \in \tilde{\Delta}_i \ \forall i$. We can write $\tilde{p}_i = p_i \times e_i \in \tilde{\Delta}_i \cap \widetilde{M}$ ($1 \leq i \leq r$). By 2.6, $\tilde{p}_1 + \cdots + \tilde{p}_r = m_{\sigma^\vee} = m \times (e_1 + \cdots + e_r)$. Hence for $i = 1, \dots, r$ we have $p_i \in \Delta_i \cap M$ with $p_1 + \cdots + p_r = m$ and $\Delta_1, \dots, \Delta_r$ is a nef-partition.

(3) $\Rightarrow$ (1): By Theorem 2.6 we know that σ is a reflexive Gorenstein cone of index r , hence also $\sigma^\vee$ is. We define $\tilde{p}_i := p_i \times e_i \in \overline{M} = M \oplus \mathbb{Z}^r$ for $i = 1, \dots, r$. Then $\tilde{p}_1, \dots, \tilde{p}_r \in \Delta_1 * \cdots * \Delta_r \cap \overline{M}$ with $\tilde{p}_1 + \cdots + \tilde{p}_r = m_{\sigma^\vee}$. Hence $\sigma^\vee$ is completely split by Cor. 2.5. $\square$

We can sum this up in a more symmetric way:

Corollary 3.7. *A reflexive Gorenstein cone σ is associated to a nef-partition if and only if both cones σ and $\sigma^\vee$ are completely split.*

A Gorenstein polytope $\tilde{\Delta}$ of index r is the Cayley polytope of a nef-partition if and only if $\tilde{\Delta}$ and $\tilde{\Delta}^$ are Cayley polytopes of length r , or equivalently, if and only if both polytopes $\tilde{\Delta}$ and $\tilde{\Delta}^*$ contain special $(r-1)$ -simplices.*

Example 3.8. In Example 2.17, the polytope $\tilde{\Delta} = \Delta_1 * \Delta_2$ is a Gorenstein polytope of index 2. It does not contain a special 1-simplex. Thus $\tilde{\Delta}^*$ is not a Cayley polytope of length 2. In other words, the Cayley cone σ associated to Δ_1, Δ_2 is completely split, while $\sigma^\vee$ is not. Hence the sum $\Delta_1 + \Delta_2$ is not a nef-partition of the reflexive polytope $\Delta_1 + \Delta_2$, because there are no lattice points $p_1 \in \Delta_1, p_2 \in \Delta_2$ such that $p_1 + p_2 = 0$.

From Cor. 2.12 we get:

Corollary 3.9. *Let $\tilde{\Delta}$ be a Gorenstein polytope such that both $\tilde{\Delta}$ and $\tilde{\Delta}^*$ are integrally closed. Then $\tilde{\Delta}$ (and also $\tilde{\Delta}^*$) is a Cayley polytope associated to a nef-partition.*

Remark 3.10. In dimension 2, all lattice polygons Δ_1, Δ_2 such that $\Delta_1 + \Delta_2$ is reflexive were listed in [ET05, Fig.4,5]. In dimension 3 and 4, all reflexive polytopes are completely classified in [KS98, KS00]. Using the software package PALP, one can find all possible centered nef-partitions

whose Minkowski sum yields a given reflexive polytope [KS04]. However, it would be desirable to have an efficient algorithm that classifies all centered nef-partitions of given length r in fixed dimension d . At the moment, the only way is to compute *all* reflexive polytopes of dimension d , which is practically impossible for $d \geq 5$.

3.2. Duality of nef-partitions.

Definition 3.11. Let P and P^* be two dual to each other d -dimensional Gorenstein polytopes of index r . Assume that there exist two special $(r-1)$ -simplices $S \subset P$ and $S' \subset P^*$. By 2.8, these simplices define Cayley structures on polytopes P and P^* . Therefore, for some lattice polytopes $\Delta_1, \dots, \Delta_r$ and $\nabla_1, \dots, \nabla_r$ we obtain

$$P = \Delta_1 * \dots * \Delta_r, \quad P^* = \nabla_1 * \dots * \nabla_r,$$

and two nef-partitions

$$\Delta = \Delta_1 + \dots + \Delta_r, \quad \nabla = \nabla_1 + \dots + \nabla_r$$

of $(d-r+1)$ -dimensional reflexive polytopes Δ and ∇ . We call these two nef-partitions *dual to each other*.

Remark 3.12. The duality of nef-partitions defined in 3.11 looks trivial, because it simply exchanges Gorenstein polytopes P with P^* and special $(r-1)$ -simplices S and S' . By 2.9, for $r=1$ this duality reduces to the polar duality between reflexive polytopes P and P^* . However, Definition 3.11 is not very convenient if we want to determine explicitly the polytopes $\nabla_1, \dots, \nabla_r$ from polytopes $\Delta_1, \dots, \Delta_r$. For this reason, we consider below another combinatorial approach to the duality of centered nef-partitions.

Proposition 3.13. *Let $\Delta \subset M_{\mathbb{R}}$ be a reflexive polytope and $0 = \text{int}(\Delta) \cap M$. Assume that $\Delta = \Delta_1 + \dots + \Delta_r$ is a centered nef-partition. Then r polytopes*

$$\nabla_i := \{y \in N_{\mathbb{R}} : \langle \Delta_j, y \rangle \geq -\delta_{ij} \ \forall j = 1, \dots, r\}, \quad i \in \{1, \dots, r\}.$$

define the dual centered nef-partition.

Proof. Consider two lattices $\overline{M} = M \oplus \mathbb{Z}e_1 \oplus \dots \oplus \mathbb{Z}e_r$, and $\overline{N} = N \oplus \mathbb{Z}e_1^* \oplus \dots \oplus \mathbb{Z}e_r^*$, where $\langle e_i, e_j^* \rangle = \delta_{ij}$. Let $\sigma \subseteq \overline{M}_{\mathbb{R}}$ be the reflexive Cayley cone associated with the lattice polytopes $\Delta_1, \dots, \Delta_r \subseteq M_{\mathbb{R}}$. Denote by $\sigma^\vee \subset \overline{N}_{\mathbb{R}}$ the dual reflexive cone. We set $\tilde{\Delta}_i = \Delta_i \times e_i$. Then $0 \times (e_1 + \dots + e_r) = m_{\sigma^\vee}$, and $0 \times (e_1^* + \dots + e_r^*) = n_{\sigma}$. Let $\tilde{\Delta}^*$ be the support of $\sigma^\vee$.

For every $i = 1, \dots, r$, we define polytopes $\tilde{\nabla}_i := \{y \in \tilde{\Delta}^* : \langle e_j, y \rangle = 0 \text{ for } j \neq i\}$. By Prop. 2.3, we have $\tilde{\Delta}^* = \text{Conv}(\tilde{\nabla}_1, \dots, \tilde{\nabla}_r)$. By Prop. 3.6, for every $i = 1, \dots, r$, we can write $\tilde{\nabla}_i$ as $\nabla_i' \times e_i^*$ for some $\nabla_i' \subseteq N_{\mathbb{R}}$. We want to show that $\nabla_i = \nabla_i'$ ($1 \leq i \leq r$). This follows from the observation that for $y \in N_{\mathbb{R}}$ and $i \in \{1, \dots, r\}$ one has

$$\langle \Delta_j, y \rangle \geq -\delta_{ij} \ \forall j = 1, \dots, r$$

if and only if

$$\langle \tilde{\Delta}_j, y \times e_i^* \rangle \geq 0 \quad \forall j = 1, \dots, r$$


(i.e., $y \times e_i^* \in \tilde{\nabla}_i \Leftrightarrow y \in \nabla_i$). Obviously, one has $0 \in \nabla_i$ ($1 \leq i \leq r$). So we obtain that $\nabla_1, \dots, \nabla_r \subseteq N_{\mathbb{R}}$ is the centered dual nef-partition. $\square$

Example 3.14. The dual centered nef-partition to the one considered in 3.2 consists of two d -dimensional standard simplices

$$\nabla_1 := \{x = (x_i) \in \mathbb{R}^d : x_1 + \dots + x_d \geq -1, x_i \leq 0 \forall i\},$$

$$\nabla_2 := \{x = (x_i) \in \mathbb{R}^d : x_1 + \dots + x_d \leq 1, x_i \geq 0 \forall i\}.$$

Example 3.15. The following picture below describes another example of two dual to each other nef-partitions in dimension 2.



In this case, the dimensions of Δ_2 and ∇_2 are not maximal.

For the remainder of this section let $\Delta_1, \dots, \Delta_r$ and $\nabla_1, \dots, \nabla_r$ be two dual to each other centered nef-partitions.

Corollary 3.16. *For any $i \in \{1, \dots, r\}$ one has $\Delta_i = 0$ if and only if $\nabla_i = 0$. In particular, the dual nef-partition of a proper nef-partition is a proper nef-partition.*

Proof. Without loss of generality, we can assume that $\Delta_1 = 0$. Then the Minkowski sum $\Delta_2 + \dots + \Delta_r = \Delta_1 + \dots + \Delta_r$ is a reflexive polytope Δ containing 0 in its interior. Hence $\{y \in N_{\mathbb{R}} : \langle x, y \rangle \geq 0 \forall x \in \Delta\} = 0$, so by the formulas in 3.13 we obtain that $\nabla_1 = 0$. By duality of nef-partitions, the condition $\nabla_1 = 0$ implies $\Delta_1 = 0$, analogously. $\square$

With the same argument we also derive from Prop. 3.13:

Corollary 3.17. $\mathbb{R}_{\geq 0}(\nabla_i) \cap \mathbb{R}_{\geq 0}(\nabla_j) = \{0\}$ for $i \neq j$.

The crucial property of the duality of nef-partitions is that it exchanges Minkowski sum and convex hull:

Proposition 3.18. *$\text{Conv}(\Delta_1, \dots, \Delta_r)$ is a reflexive polytope which is dual to $\nabla_1 + \dots + \nabla_r$. Moreover, the two lattice polytopes $\Delta_1 * \dots * \Delta_r$ and $\text{Conv}(\Delta_1, \dots, \Delta_r)$ have the same h^* -polynomial.*

Proof. The statements follow immediately from Theorem 2.16, because $\text{Conv}(\Delta_1, \dots, \Delta_r)$ is the projection of $\Delta_1 * \dots * \Delta_r$ along $\text{aff}(e_1, \dots, e_r)$. $\square$

Hence, we have for $\Delta = \Delta_1 + \dots + \Delta_r$ and $\nabla = \nabla_1 + \dots + \nabla_r$:

$$\Delta^* = \text{Conv}(\nabla_1, \dots, \nabla_r) \text{ and } \nabla^* = \text{Conv}(\Delta_1, \dots, \Delta_r).$$

Now, we show that in order to determine the dual nef-partition it suffices to dualize the Minkowski sum Δ , and to choose the vertices of $\nabla_1, \dots, \nabla_r$ among the vertices of Δ^* . For this we take a closer look at boundary of the lattice polytopes $\nabla_1, \dots, \nabla_r$:

Proposition 3.19. *We have for $i = 1, \dots, r$:*

$$\nabla_i \cap \partial\Delta^* = \{v \in \Delta^* : \min_{u \in \Delta_i} \langle u, v \rangle = -1\}.$$

Any face F of ∇_i , with $0 \notin F$, is a face of $\Delta^ = \text{Conv}(\nabla_1, \dots, \nabla_r)$ and contained in a face of $\nabla = \nabla_1 + \dots + \nabla_r$.*

Proof. Let $v \in \partial\Delta^*$. Then

$$\left(\min_{u \in \Delta_1} \langle u, v \rangle\right) + \dots + \left(\min_{u \in \Delta_r} \langle u, v \rangle\right) = \min_{u \in \Delta} \langle u, v \rangle = -1,$$

and these are all numbers in $[0, -1]$. From Prop. 3.13 the equation follows.

Now, let F be a facet of ∇_i , with $0 \notin F$. Let v be a point in the relative interior of F . By Prop. 3.13 we have $\min_{u \in \Delta_i} \langle u, v \rangle = -1$, so there is a vertex u of Δ_i such that $\langle u, v \rangle = -1$. Therefore $F = \{y_i \in \nabla_i : \langle u, y_i \rangle = -1\}$, and F is also contained in the face $\{y \in \nabla_1 + \dots + \nabla_r : \langle u, y \rangle = -1\}$ of $\nabla_1 + \dots + \nabla_r$.

We show that $F = \{y \in \Delta^* : \langle u, y \rangle = -1\}$. Let $y \in \Delta^*$ with $\langle u, y \rangle = -1$. There is a convex combination $y = \sum_{j=1}^r \lambda_j v_j$, with $\lambda_j \geq 0$, $\sum_{j=1}^r \lambda_j = 1$, for $v_j \in \nabla_j$ ($j = 1, \dots, r$). Hence from $-1 = \langle u, y \rangle$ we get $-1 \geq \lambda_i \langle u, v_i \rangle \geq -\lambda_i$, because of Prop. 3.13, thus $\lambda_i = 1$, so $y = v_i \in \nabla_i$. This shows $y \in F$. $\square$

Remark 3.20. The reader further interested in faces of nef-partitions and their dualities is encouraged to consult [HZ05, Prop.2.3] and [Gro05, 2.7-2.9].

As mentioned above, here is a direct way to determine the vertices of the dual nef-partition:

Corollary 3.21. *We have for $i = 1, \dots, r$:*

$$\mathcal{V}(\nabla_i) \setminus \{0\} = \{v \in \mathcal{V}(\Delta^*) : \min_{u \in \Delta_i} \langle u, v \rangle = -1\}.$$

Remark 3.22. The previous statement implies that any non-zero vertex of ∇_i ($i = 1, \dots, r$) is a vertex of $\text{Conv}(\nabla_1, \dots, \nabla_r)$. It is not difficult to prove, that this also holds under the weaker assumption that $\nabla_1, \dots, \nabla_r$ are lattice polytopes, each containing 0, such that $\nabla_1 + \dots + \nabla_r$ has no interior lattice point except 0.

Not only vertices but also lattice points of $\Delta^* = \text{Conv}(\nabla_1, \dots, \nabla_r)$ necessarily belong to one of the lattice polytopes $\nabla_1, \dots, \nabla_r$:

Corollary 3.23. *Any non-zero lattice point of Δ^* is contained for some $i \in \{1, \dots, r\}$ in a face of ∇_i that does not contain the origin. In particular $|\Delta^* \cap N| = |\nabla_1 \cap N| + \dots + |\nabla_r \cap N| - r + 1$.*

Proof. Let $v \in \Delta^* \cap N$, $v \neq 0$. There has to be some $u \in \mathcal{V}(\Delta_i)$ ($i \in \{1, \dots, r\}$) with $\langle u, v \rangle < 0$, thus $\langle u, v \rangle = -1$. Now use Prop. 3.19. $\square$

4. THE E_{st} -FUNCTION OF A GORENSTEIN POLYTOPE

Borisov and the first author gave in [BB96b] an explicit formula for the generating function of stringy Hodge numbers of Calabi-Yau complete intersections given by a nef-partition (the stringy Hodge numbers were introduced in [BD96, Bat98]). We remark that this formula can be used for arbitrary reflexive Gorenstein cones or for arbitrary Gorenstein polytopes. However, it is not clear a priori whether the corresponding E_{st} -function is a polynomial. We expect that it is always the case.

In this section, we review the above formula using $\tilde{S}$ -polynomials and ideas from [BM03]. Our goals are to introduce the E_{st} -function of a Gorenstein polytope, and to show some of its properties, and to formulate related open questions and conjectures.

4.1. The $\tilde{S}$ -polynomial of a lattice polytope. We start by recalling the definition of the g - and h -polynomials, defined by Stanley for general Eulerian posets in [Sta87] (see also [BM03, Def.5.3]):

Definition 4.1. Let $\mathcal{P}$ be an Eulerian poset of rank d , $\hat{0}$ its minimal element, $\hat{1}$ its maximal element. Then the polynomials $g_{\mathcal{P}}(t), h_{\mathcal{P}}(t) \in \mathbb{Z}[t]$ are defined recursively as follows. If $d = 0$ then we set $g_{\mathcal{P}}(t) := h_{\mathcal{P}}(t) := 1$. If $d > 0$ then the h -polynomial is defined as

$$h_{\mathcal{P}}(t) = \sum_{\hat{0} < x \leq \hat{1}} (t-1)^{\text{rk}(x)-1} g_{[x, \hat{1}]}(t).$$

For $d > 0$, we define the g -polynomial $g_{\mathcal{P}}(t) = \sum_i g_i t^i$ as a polynomial of degree $< d/2$ whose coefficients g_i (for $0 \leq i < d/2$) are equal to the coefficients c_i of the polynomial

$$(1-t)h_{\mathcal{P}}(t) = \sum_i c_i t^i,$$

i.e. , $g_i = h_i - h_{i-1}$ for $0 \leq i < d/2$.

A recent account on this important combinatorial notion can be found in [Bra05]. Here we need only the following well-known properties:

Remark 4.2. The polynomials $g_{\mathcal{P}}(t)$ and $h_{\mathcal{P}}(t)$ have the following properties:

- (1) every coefficient of $g_{\mathcal{P}}(t)$ is a non-negative integer;
- (2) $g_{\mathcal{P}}(t) = 1$ if and only if the poset $\mathcal{P}$ is boolean, i.e., it is isomorphic to the poset of all faces (including the empty one) of a $(d-1)$ -dimensional simplex;
- (3) the polynomial g is multiplicative, i.e., $g_{\mathcal{P} \times \mathcal{P}'}(t) = g_{\mathcal{P}}(t) \cdot g_{\mathcal{P}'}(t)$.

Combinatorial data contained in the g -polynomial will be combined with the enumeration of lattice points via the h^* -polynomial in the following definition [BM03, Def.5.3]:

Definition 4.3. Let P be a lattice polytope. We define the $\tilde{S}$ -polynomial:

$$\tilde{S}(P, t) := \sum_{\emptyset \leq F \leq P} (-1)^{\dim(P) - \dim(F)} h_F^*(t) g_{[F, P]}(t),$$

where $h_F^*(t)$ is the h^* -polynomial of the lattice polytope F (with convention $h_\emptyset^*(t) := 1$, see Def. 2.14) and $g_{[F, P]}(t)$ is the g -polynomial of the poset $[F, P]$ of faces between F and P . Note that $\tilde{S}(\emptyset, t) = 1$. However, $\tilde{S}(P, t) = 0$, if $\dim(P) = 0$.

Remark 4.4. The $\tilde{S}$ -polynomial of a lattice polytope P has the following properties:

- (1) all coefficients of $\tilde{S}(P, t)$ are non-negative integers (this follows from [BM03, Prop.5.5], where it is shown that $\tilde{S}(P, t)$ is the Hilbert function of a graded vector space).
- (2) one has the reciprocity-law (see [BM03, Remark 5.4]):

$$\tilde{S}(P, t) = t^{\dim(P)+1} \tilde{S}(P, 1/t).$$

Here is another nice property of the $\tilde{S}$ -polynomial:

Lemma 4.5. Let P be a lattice polytope. If P is a lattice pyramid, then $\tilde{S}(P, t) = 0$.

Proof. Let P be a lattice pyramid over the facet P' with apex v , i.e., $h_{P'}^*(t) = h_P^*(t)$. Any face F of P , that is not contained in P' , is a pyramid over a face F' of P' with apex v . Hence, $h_F^*(t) = h_{F'}^*(t)$ and $g_{[F, P]}(t) = g_{[F', P']}(t)$. On the other hand, if F is a face of P' , then $g_{[F, P]}(t) = g_{[F, P']}(t)$, since $[F, P]$ is a product of $[F, P']$ and an interval, and g is multiplicative. This yields:

$$\tilde{S}(P, t) = \tilde{S}(P', t) + (-1)^{\dim(P) - \dim(P')} \tilde{S}(P', t) = 0.$$

□

In the case of a simplex, the $\tilde{S}$ -polynomial is easy to calculate:

Proposition 4.6. Let $P \subseteq M_{\mathbb{R}}$ be a lattice simplex. Let $\tilde{P} := P \times 1 \subseteq \overline{M}_{\mathbb{R}} = M_{\mathbb{R}} \oplus \mathbb{R}$. We denote by Π the parallelepiped spanned by the vertices of $\tilde{P}$. Then

$$\tilde{S}(P, t) = \sum_{x \in \text{int}(\Pi) \cap M} t^{x_{d+1}}.$$

The degree of $\tilde{S}(P, t)$ equals at most the degree of $h_P^*(t)$.

Proof. For a simplex $\tilde{P}$, the h^* -polynomial can be calculated by counting the lattice points in the parallelepiped $\Pi(\tilde{P}) := \{\sum_{i=0}^d \lambda_i v_i : 0 \leq \lambda_i < 1\}$, where $v_0, \dots, v_d$ are the vertices of $\tilde{P}$:

$$h_{\tilde{P}}^*(t) = \sum_{x \in \Pi(\tilde{P}) \cap M} t^{x_{d+1}}.$$

Since the g -polynomial of a simplex equals 1, we get

$$\tilde{S}(P, t) = \sum_{\emptyset \leq F \leq \tilde{P}} (-1)^{\dim(\tilde{P}) - \dim(F)} h_F^*(t).$$

Now, for $x \in \Pi(\tilde{P})$, we denote by F_x the smallest face of $\tilde{P}$ such that $x \in \mathbb{R}_{\geq 0}(F_x)$. Hence,

$$\tilde{S}(P, t) = \sum_{x \in \Pi(\tilde{P}) \cap M} \left(\sum_{F_x \leq F \leq \tilde{P}} (-1)^{\dim(\tilde{P}) - \dim(F)} \right) t^{x_{d+1}}.$$

The Euler-Poincaré-formula implies that the right side vanishes, whenever $F_x \neq \tilde{P}$. From this the equation follows. $\square$

In general, the degree of the $\tilde{S}$ -polynomial of a lattice simplex is not equal to the degree of the h^* -polynomial. This follows immediately from Lemma 4.5.

Example 4.7. For any lattice polytope $P \subseteq M_{\mathbb{R}}$ we denote by $l^*(P)$ (resp. by $l(P)$) the number of M -lattice points in the relative interior of P (resp. in $P \cap M$).

Now, let P be a 2-dimensional lattice polytope with k vertices (a lattice k -gon). In this case, we calculate from the definition of the h - and g -polynomial, $h_{[\emptyset, P]}(t) = k + k(t-1) + (t-1)^2 = 1 + (k-2)t + t^2$, thus $g_{[\emptyset, P]}(t) = 1 + (k-3)t$. Then

$$\begin{aligned} \tilde{S}(P, t) &= h_P^*(t) - \left(\sum_{\dim \Gamma=1} h^*(\Gamma) \right) + k - (1 + (k-3)t) = \\ &= 1 + (l(P) - 3)t + l^*(P)t^2 - \left(\sum_{\dim \Gamma=1} 1 + (l^*(\Gamma) - 1)t \right) + k - (1 + (k-3)t) = \\ &= l^*(P)(t + t^2), \end{aligned}$$

since

$$l(P) = l^*(P) + \sum_{\dim \Gamma=1} l^*(\Gamma) + k.$$

More generally, if P is a d -dimensional lattice polytope and $l^*(P) \neq 0$, then $\tilde{S}(P, t)$ and $h_P^*(t)$ have the same leading term $l^*(P)t^d$, because

$$\deg(h_F^*(t) g_{[F, P]}(t)) \leq d - 1$$

for any proper face $F \subset P$, since $\deg(h_F^*(t)) \leq \dim(F)$ and $\deg(g_{[F, P]}(t)) < (\text{rank}[F, P])/2 = (d - \dim(F))/2$.

4.2. E_{st} -function of a Gorenstein polytope.

Definition 4.8. Let Δ be a d -dimensional Gorenstein polytope of index r . We define the following function of two independent variables u and v :

$$E_{\text{st}}(\Delta; u, v) := (uv)^{-r} \sum_{\emptyset \leq F \leq \Delta} (-u)^{\dim(F)+1} \tilde{S}(F, u^{-1}v) \tilde{S}(F^*, uv),$$

where F^* is the face of Δ^* dual to F . We call it the *stringy E -function* (or E_{st} -function) of Δ . The integer

$$\text{CY-dim}(\Delta) := d + 1 - (2r)$$

we call the *Calabi-Yau dimension* (CY-dimension) of Δ .

Remark 4.9.

- (1) The E_{st} -function is symmetric

$$E_{\text{st}}(\Delta; u, v) = E_{\text{st}}(\Delta; v, u)$$

and it satisfies the Poincaré duality

$$(uv)^{\text{CY-dim}(\Delta)} E_{\text{st}}(\Delta; u^{-1}, v^{-1}) = E_{\text{st}}(\Delta; u, v);$$

- (2) The E_{st} -function satisfies the following reciprocity-law:

$$E_{\text{st}}(\Delta; u, v) = (-u)^{\text{CY-dim}(\Delta)} E_{\text{st}}(\Delta^*; u^{-1}, v).$$

Note, that we have by definition $\text{CY-dim}(\Delta) = \text{CY-dim}(\Delta^*)$.

- (3) Let $\deg(\Delta)$ be the degree of the h^* -polynomial of Δ . Then we have $\text{CY-dim}(\Delta) = \deg(\Delta) - r$, cf. [Bat07].
- (4) The CY-dimension can be arbitrarily negative: If Δ is a lattice pyramid over Δ' , then $\text{CY-dim}(\Delta) = \text{CY-dim}(\Delta') - 1$, since the index of Δ is 1 plus the index of Δ' .

Here is our conjecture on the E_{st} -function:

Conjecture 4.10. Let Δ be a Gorenstein polytope of CY-dimension n . Then

$$E := E_{\text{st}}(\Delta; u, v) = \sum_{p, q} (-1)^{p+q} h_{\Delta}^{p, q} u^p v^q$$

is a polynomial in u, v of degree $2n$ with nonnegative integral coefficients $h_{\Delta}^{p, q}$ (in particular, one has $E = 0$ if $n < 0$, and E is constant if $n = 0$). Moreover, for any $n \geq 1$ the polynomial E satisfies the following two conditions

$$E_{\text{st}}(\Delta; u, 0) = (-u)^n E_{\text{st}}(\Delta; u^{-1}, 0)$$

and

$$\frac{d^2}{du^2} E_{\text{st}}(\Delta; u, 1)|_{u=1} = \frac{n(3n-5)}{12} E_{\text{st}}(\Delta; 1, 1).$$

These two conditions together with (1) and (2) from 4.9 imply:

- (1) If $n = 1$, then for some integer k one has

$$E_{\text{st}}(\Delta; u, v) = k(1-u)(1-v).$$

(2) If $n = 2$, then for some integers k, l one has

$$E_{\text{st}}(\Delta; u, v) = k(1 + u^2)(1 + v^2) - 2l(u + v)(1 + uv) + (20k - 16l)uv.$$

In particular, the number $E_{\text{st}}(\Delta; 1, 1) = 24(k - l)$ is always divisible by 24.

Remark 4.11. This conjecture is motivated by the expectation that the function $E_{\text{st}}(\Delta; u, v)$ “looks” as a generating function for stringy Hodge numbers of a disjoint union of manifolds of dimension $\text{CY-dim}(\Delta)$ with trivial canonical class. The second condition was proved for stringy Hodge numbers in a more general form in [Bat00]. In the case $\text{CY-dim}(\Delta) = 0$ we expect a finite union of points. In the case $\text{CY-dim}(\Delta) = 1$, it should be a union of elliptic curves. In the case $\text{CY-dim}(\Delta) = 2$, one expects Hodge numbers of abelian surfaces and $K3$ -surfaces.

In all examples that the authors know the leading coefficient k is either zero, or a power of 2 (see also 4.22).

Example 4.12. Let S_d be a standard unimodular simplex of dimension $d := 2r - 1$ ($r \in \mathbb{N}$). We calculate the E_{st} -function of the Gorenstein simplex $\Delta := 2S_d$ which has index r . We have $\text{CY-dim}(\Delta) = 0$. By Lemma 1.16 the dual Gorenstein polytope Δ^* is a Gorenstein simplex with only unimodular facets. Hence, Lemma 4.5 yields:

$$E_{\text{st}}(\Delta; u, v) = (uv)^{-r} \left(\tilde{S}(\Delta^*, uv) + (-u)^{2r} \tilde{S}(\Delta, u^{-1}v) \right).$$

We claim:

$$(*) \quad \tilde{S}(\Delta, t) = t^r = \tilde{S}(\Delta^*, t).$$

Combining these two equations we get:

$$E_{\text{st}}(\Delta; u, v) = 2.$$

We show $(*)$, by proving that $\tilde{S}(\Delta, t) \in \{0, t^r\}$ for any Gorenstein simplex Δ of index r and with $\text{CY-dim}(\Delta) = 0$. By Prop. 4.6 and its proof, to determine $\tilde{S}(\Delta, t)$ we have to compute the sum $\sum_x t^{x_{d+1}}$ over the lattice points x in the interior of the parallelepiped spanned by vertices of Δ . On the other hand, in order to determine $h_{\Delta}^*(t)$ one computes the sum $\sum_x t^{x_{d+1}}$ over *all* lattice points x in the parallelepiped. By Remark 4.9(3), the degree of h_{Δ}^* equals the index r . Hence, there is only one interior lattice point x in the parallelepiped such that $x_{d+1} = r$.

In this case, x is the unique interior lattice point in $r\Delta = (2r)S_d$ which is given as $\sum_{v \in \mathcal{V}(\Delta)} \frac{1}{2}v$. Hence $\tilde{S}(\Delta, t) = t^r$. In the same way, this holds for Δ^* . This proves $(*)$.

From Remark 4.9(4) we see that using lattice pyramids we may construct Gorenstein polytopes of negative CY-dimension. In this case, the E_{st} -function should vanish. However, this vanishing holds for any lattice pyramid (even of positive CY-dimension):

Proposition 4.13. *Let Δ be a Gorenstein polytope. If Δ is a lattice pyramid, then $E_{\text{st}}(\Delta; u, v) = 0$.*

Proof. Let Δ be a lattice pyramid over a facet Δ' with apex v . Then also Δ^* is a lattice pyramid over the facet v^* with apex $(\Delta')^*$, for instance, use Prop. 2.3. Thus, for any face F of Δ , F or F^* is a lattice pyramid. Hence, $\tilde{S}(F, u^{-1}v) \tilde{S}(F^*, uv) = 0$ by Lemma 4.5. From this the statement follows. $\square$

4.3. The B -polynomial. Originally, in [BB96b] so-called B -polynomials were used to calculate Hodge numbers, see also [KRS03, Remark 5.8] and [Haa00]. In [BM03, Lemma 11.4] Borisov and Mavlyutov gave the following definition in terms of g -polynomials:

Definition 4.14. Let $\mathcal{P}$ be an Eulerian poset of rank d , with minimal element $\hat{0}$ and maximal element $\hat{1}$. Then

$$B(\mathcal{P}; u, v) := \sum_{\hat{0} \leq x \leq \hat{1}} (-u)^{\text{rk}(\hat{1}) - \text{rk}(x)} g_{[x, \hat{1}]^*}(u^{-1}v) g_{[\hat{0}, x]}(uv).$$

To see, how the B -polynomial naturally appears, we separately collect all g -polynomials in the definition of the E_{st} -function into one polynomial in two variables: Let Δ be a Gorenstein polytope of index r . Then $E_{\text{st}}(\Delta; u, v)$ equals

$$(uv)^{-r} \sum_{F, F'} (-1)^{\dim((F')^*) + 1} u^{\dim(F) + 1} h_F^*(u^{-1}v) h_{F'}^*(uv) B([F, (F')^*]; u, v),$$

where the sum is over all pairs of faces $\emptyset \leq F \leq \Delta$, $\emptyset \leq F' \leq F^*$.

There is an important identity on g -polynomials, called Stanley's convolution, cf. [Sta92, Cor.8.3], that can be stated in the following form, see [BM03, Lemma 11.2],

Lemma 4.15. *$B([F, (F')^*]; 1, v)$ equals 0, if $F \neq (F')^*$, and 1, otherwise.*

As a corollary we get

$$E_{\text{st}}(\Delta; 1, v) = v^{-r} \sum_{\emptyset \leq F \leq \Delta} (-1)^{\dim(F) + 1} h_F^*(v) h_{F^*}^*(v),$$

$$(**) \quad E_{\text{st}}(\Delta; 1, 1) = \sum_{\emptyset \leq F \leq \Delta} (-1)^{\dim(F) + 1} \text{Vol}(F) \text{Vol}(F^*),$$

where Vol is the lattice volume (with convention $\text{Vol}(\emptyset) := 1$).

Remark 4.16. When Δ is a three-dimensional reflexive polytope, Theorem 4.17 below yields that $E_{\text{st}}(\Delta; 1, 1)$ equals the topological Euler-characteristic of a $K3$ surface. Hence, $E_{\text{st}}(\Delta; 1, 1) = 24$. However, there is still no direct combinatorial proof for this equation, cf. [Haa05].

4.4. Stringy Hodge numbers. The main motivation for the definition of the E_{st} -function and the Calabi-Yau dimension, and for the related conjectures stems from the following theorem:

Theorem 4.17. [BM03, Thm.7.2] *Let σ be a $(d+r)$ -dimensional reflexive Gorenstein cone of index r with support $\tilde{\Delta} \subseteq \tilde{M}_{\mathbb{R}}$, where $\tilde{\Delta}$ is a Cayley polytope of r lattice polytopes. Let Y be a generic complete intersection Calabi-Yau defined by these r equations, hence $\dim(Y) = d - r = \text{CY-dim}(\tilde{\Delta})$. Then*

$$E_{\text{st}}(\tilde{\Delta}; u, v) = E_{\text{st}}(Y; u, v) = \sum_{p,q} (-1)^{p+q} h_{\text{st}}^{p,q}(Y) u^p v^q$$

is a polynomial in u, v , called E_{st} -polynomial of Y , where the coefficients $h_{\text{st}}^{p,q}$ are the so-called stringy Hodge numbers of Y , in the sense of [BD96, Bat98].

This is a simplified version of the main result in [BB96b]. Note, that the combinatorial identity (**) gives the topological Euler-characteristic of the Calabi-Yau Y . From this theorem we can derive the following result using algebraic geometry (for $n = 2$, Y is a $K3$ -surface) and the reciprocity-law (Remark 4.9(2)):

Corollary 4.18. *Conjecture 4.10 holds, if at least one of the two Gorenstein polytopes Δ or Δ^* of index r is a Cayley polytope of length r .*

Example 4.19. Here is an example for a Gorenstein polytope Δ that does not satisfy the assumption of the previous corollary. Let $\Delta_1 := 2S_3$, where S_3 is the unimodular 3-simplex, and let σ_1 be the reflexive Gorenstein cone of index two with support Δ_1 . We also define $\Delta_2 := \Delta_1^*$, and $\sigma_2 := \sigma_1^\vee$. Then by Lemma 4.20 below, the support Δ of the direct sum $\sigma_1 \oplus \sigma_2$ is a 7-dimensional Gorenstein polytope of index 4. The dual Gorenstein polytope Δ^* is the support of $\sigma_2 \oplus \sigma_1$. Since Δ_2 does not contain a special 1-simplex, both Δ and Δ^* do not contain special 3-simplices, hence by Prop. 2.8 both Gorenstein polytopes of index 4 are not Cayley polytopes of length 4.

The proof of the following lemma is left to the reader.

Lemma 4.20. *Let $\sigma_1 \subseteq \overline{M}_{\mathbb{R}}$, $\sigma_2 \subseteq \overline{M}'_{\mathbb{R}}$ be two reflexive Gorenstein cones of indices r_1 , respectively r_2 . Then the direct sum $\sigma_1 \oplus \sigma_2 \subseteq (\overline{M} \oplus_{\mathbb{Z}} \overline{M}')_{\mathbb{R}}$ is a reflexive Gorenstein cone of index $r_1 + r_2$, with dual cone $\sigma_1^\vee \oplus \sigma_2^\vee$. Moreover, the support of $\sigma_1 \oplus \sigma_2$ contains a special $(r_1 + r_2 - 1)$ -simplex if and only if the support of σ_1 contains a special $(r_1 - 1)$ -simplex and the support of σ_2 contains a special $(r_2 - 1)$ -simplex.*

If in Theorem 4.17 the cone σ is associated to a nef-partition, then we get from Remark 4.9(2) for the E_{st} -polynomial of Y

$$E_{\text{st}}(Y; u, v) = (-u)^{d-r} E_{\text{st}}(Y^*; u^{-1}, v),$$

where Y^* is a $(d - r)$ -dimensional complete intersection Calabi-Yau defined by the dual nef-partition. In particular, stringy Hodge numbers satisfy the following mirror symmetry property:

$$h_{\text{st}}^{p,q}(Y) = h_{\text{st}}^{d-r-p,q}(Y^*) \quad \text{for } 0 \leq p, q \leq d - r.$$

4.5. Boundedness of Calabi-Yau n -folds. It is still an open question whether there is only a finite number of topologically different Calabi-Yau n -folds ($n \geq 3$). We can look at this question from a combinatorial point of view and ask another one:

Question 4.21. Let n be fixed. Is there up to scalar only a finite number of polynomials in u, v that occur as E_{st} -polynomials of Gorenstein polytopes Δ such that $\text{CY-dim}(\Delta)$ equals a fixed number n ?

Remark 4.22. In particular, a positive answer would imply that the set of stringy Hodge numbers associated to irreducible Calabi-Yau complete intersections Y defined by a nef-partition is bounded. This implication follows from the fact that by Poincaré duality the leading coefficient k of $E_{\text{st}}(Y; u, v)$ equals the constant coefficient which is 1, because $k = h^{0,0}(Y) = h^0(O_Y)$ is the number of irreducible components of Y . We observe that in general the number k can be arbitrary large powers of 2. Indeed, let $\Delta_1, \dots, \Delta_r$ be a nef-partition of length r and let E be E_{st} -polynomial of the corresponding Calabi-Yau complete intersection Y of dimension $d - r$. Then $r + 1$ polytopes

$$(\Delta_1 \times 0), \dots, (\Delta_r \times 0), (0 \times [-1, 1]) \subseteq M_{\mathbb{R}} \oplus \mathbb{R}$$

define a reducible nef-partition of length $r + 1$ of the same CY-dimension $d - r$. The corresponding Calabi-Yau complete intersection Y' is a product of Y with two points. Therefore E_{st} -polynomial of Y' equals $2E$. By repeating the same procedure, we get E_{st} -polynomials with arbitrary large leading coefficient k .

Remark 4.23. There are various finiteness results on toric Fano varieties [Bat82, BorBor93]. However these results are not sufficient to prove finiteness of E_{st} -polynomials of Gorenstein polytopes of fixed Calabi-Yau dimension. We are still optimistic and want to illustrate our hope by an example of d -dimensional Gorenstein simplices $S(\omega)$ of index r defined as

$$S(\omega) := \{(x_0, x_1, \dots, x_d) \in \mathbb{R}_{\geq 0}^{d+1} : w_0 x_0 + w_1 x_1 + \dots + w_d x_d = w\},$$

where $w, w_0, w_1, \dots, w_d$ are positive integers having the property $w_i | w \ \forall i$. Let $k_i := w/w_i$ ($0 \leq i \leq d$). The fact that $S(\omega)$ is a Gorenstein simplex of index r is equivalent to the equation

$$\frac{1}{k_0} + \frac{1}{k_1} + \dots + \frac{1}{k_d} = r.$$

If $k_i = 1$ for some i , then $S(\omega)$ is a lattice pyramid over a $(d - 1)$ -dimensional simplex and, by 4.13, the E_{st} -polynomial of $S(\omega)$ is 0. Therefore it is enough

to consider only the case when $k_i \geq 2$ ($0 \leq i \leq d$). Let s be the number of indices i such that $k_i \geq 3$. Then we get from the above equation

$$r = \frac{1}{k_0} + \frac{1}{k_1} + \cdots + \frac{1}{k_d} \leq \frac{d+1-s}{2} + \frac{s}{3},$$

or $3(d+1-2r) \geq s$. By definition

$$\text{CY-dim}(S(\omega)) = d+1-2r.$$

So we obtain $s \leq 3\text{CY-dim}(S(\omega))$. This implies that if $S(\omega)$ has a fixed Calabi-Yau dimension $n = d+1-2r$, then there exist only finitely many possibilities for integers k_i which are different from 2. In order to compute $E_{\text{st}}(S(\omega); u, v)$ we observe that the simplex $S(\omega)$ determines a Landau-Ginzburg model with LG-potential of Fermat type:

$$F(z) := z_0^{k_0} + z_1^{k_1} + \cdots + z_d^{k_d}.$$

The corresponding formula for the E_{st} -polynomial is expected to coincide with the well-known formula of Vafa [Va89]. The latter one does not change if one adds any quadratic terms to $F(z)$. Therefore we come to finitely many E_{st} -polynomials for a fixed value of $\text{CY-dim}(S(\omega))$.


5. NEF-PARTITIONS AND HOMOLOGICAL MIRROR SYMMETRY

Let P and P^* be dual to each other Gorenstein polytopes of index $r > 1$. Example 2.10 shows that P and P^* may contain many different special $(r-1)$ -simplices. Consider the case when a Gorenstein polytope P contains two different special $(r-1)$ -simplices S' and S'' . Assume that the dual Gorenstein polytope P^* also contains a special $(s-1)$ -simplex S . Then the special $(r-1)$ -simplex S defines a Cayley structure on P , i.e., some lattice polytopes $\Delta_1, \dots, \Delta_r$ such that $P \cong \Delta_1 * \cdots * \Delta_r$. Now two different special $(r-1)$ -simplices $S', S'' \subset P$ define two collections of lattice points $p'_i, p''_i \in \Delta_i$ ($1 \leq i \leq r$) such that

$$m = p'_1 + \cdots + p'_r = p''_1 + \cdots + p''_r$$

is the unique interior lattice point in the reflexive polytope $\Delta_1 + \cdots + \Delta_r$. We remark that two Cayley structures of the dual Gorenstein polytope P^* corresponding to S' and S'' may define very different polytopes $\nabla_1, \dots, \nabla_r$.

Example 5.1. We consider below a picture of two centered nef-partitions corresponding to polytopes Δ'_1, Δ'_2 and Δ''_1, Δ''_2 which are the same up to a affine translation, i.e. $\Delta''_1 = \Delta'_1 - (0, 1)$ and $\Delta''_2 = \Delta'_2 - (0, -1)$:



We see that the corresponding polytopes ∇'_1, ∇'_2 and ∇''_1, ∇''_2 are not isomorphic via affine translations by lattice vectors.

It is easy to see that, if two nef-partitions consist of collections of lattice polytopes $\{\Delta'_1, \dots, \Delta'_r\}$ and $\{\Delta''_1, \dots, \Delta''_r\}$ such that $\Delta''_i = \Delta'_i + q_i$ for $q_i \in M$ ($1 \leq i \leq r$), then the corresponding generic Calabi-Yau complete intersections $\overline{X}_{\Delta'_i}$ and $\overline{X}_{\Delta''_i}$ are isomorphic. However, there is no natural isomorphism between the generic mirror Calabi-Yau complete intersections $\overline{X}_{\nabla'_i}$ and $\overline{X}_{\nabla''_i}$. Using the formula for E_{st} -polynomial of P^* , we can only say that the stringy Hodge numbers of $\overline{X}_{\nabla'_i}$ and $\overline{X}_{\nabla''_i}$ are the same.

Question 5.2. *Are the mirror Calabi-Yau complete intersections $\overline{X}_{\nabla'_i}$ and $\overline{X}_{\nabla''_i}$ birationally isomorphic?*

Even if in general the answer to this question will be negative, we can use the approach of the Homological Mirror Symmetry and formulate the following conjecture:

Conjecture 5.3. *There exists a equivalence (of Fourier-Mukai type) between the derived category of coherent sheaves on two Calabi-Yau complete intersections $\overline{X}_{\nabla'_i}$ and $\overline{X}_{\nabla''_i}$.*

Remark 5.4. This conjecture needs some comments, because Calabi-Yau varieties $\overline{X}_{\nabla'_i}$ and $\overline{X}_{\nabla''_i}$ are usually singular. However, using toric methods, one can always construct their partial crepant desingularizations $\widehat{X}_{\nabla'_i}$ and $\widehat{X}_{\nabla''_i}$ which are Calabi-Yau varieties having at worst terminal abelian quotient singularities. As in the toroidal case considered by Kawamata in [Ka05] the equivalence class of stacky derived categories of $\widehat{X}_{\nabla'_i}$ and $\widehat{X}_{\nabla''_i}$ should be independent of the choice of the partial crepant desingularizations. So we expect the existence of a Fourier-Mukai transform which establishes an equivalence between the stacky derived categories of $\widehat{X}_{\nabla'_i}$ and $\widehat{X}_{\nabla''_i}$.

6. OPERATIONS WITH NEF-PARTITIONS

In this concluding section we give as an addendum to Section 3 a concise survey on how to modify, project and decompose nef-partitions.

Throughout, let $\Delta_1, \dots, \Delta_r \subseteq M_{\mathbb{R}}$ and $\nabla_1, \dots, \nabla_r \subseteq N_{\mathbb{R}}$ be two dual to each other centered proper nef-partitions.

6.1. Collecting. By collecting lattice polytopes in a nef-partition we obtain another nef-partition. For this, we simply replace some lattice polytopes in a nef-partition by their Minkowski sum. This yields a shorter nef-partition, which is by Cor. 3.21 dual to the nef-partition we get by replacing the respective lattice polytopes in the dual nef-partition by their convex hull:

Definition 6.1. For $I \subseteq \{1, \dots, r\}$ we define $\nabla_I := \text{Conv}(\nabla_i : i \in I)$ and $\nabla^I := \sum_{i \in I} \nabla_i$. In the same way we define Δ_I and Δ^I .

Proposition 6.2. Let $I_1, \dots, I_l$ be a partition of $\{1, \dots, r\}$ into non-empty subsets. Then $\Delta^{I_1}, \dots, \Delta^{I_l}$ and $\nabla_{I_1}, \dots, \nabla_{I_l}$ are dual nef-partitions.

From Prop. 3.13 we get:

Corollary 6.3. For $I \subseteq \{1, \dots, r\}$ we have

$$\begin{aligned} \Delta^I &= \{x \in \Delta_1 + \dots + \Delta_r : \langle x, \nabla_j \rangle \geq 0 \ \forall j \notin I\}, \\ \nabla_I &= \{y \in \text{Conv}(\nabla_1, \dots, \nabla_r) : \langle \Delta_j, y \rangle \geq 0 \ \forall j \notin I\}. \end{aligned}$$

Remark 6.4. Let us explain how Prop. 6.2 relates to the results of Section 2: More generally, let $\Delta_1, \dots, \Delta_r$ be arbitrary lattice polytopes with $\Delta_1 + \dots + \Delta_r$ reflexive. Then by Theorem 2.6, $P := \Delta_1 * \dots * \Delta_r$ is a Gorenstein Cayley polytope of index r . The dual Gorenstein polytope P^* contains a special $(r-1)$ -simplex with vertices $v_1, \dots, v_r$ (see Prop. 2.8). Theorem 2.16 shows that projecting P^* along the affine hull of this special simplex yields the reflexive polytope $(\Delta_1 + \dots + \Delta_r)^*$.

Now, let $I_1, \dots, I_l$ be a partition of $\{1, \dots, r\}$ into non-empty subsets. We may assume $I_1 = \{1, \dots, r'\} \subseteq \{1, \dots, r\}$. In the same way as in the proof of Theorem 2.6 and Theorem 2.16, projecting P^* along $\text{aff}(v_1, \dots, v_{r'})$ yields a Gorenstein polytope dual to the Gorenstein polytope $\Delta^{I_1} * \Delta_{r'+1} * \dots * \Delta_r$ of index $r+1-r'$. After l such projection steps, we finally get a Gorenstein polytope of index l dual to $\Delta^{I_1} * \Delta^{I_2} * \dots * \Delta^{I_l}$.

In particular, when $\Delta_1, \dots, \Delta_r$ and $\nabla_1, \dots, \nabla_r$ are dual to each other centered nef-partitions, we see that $P^* = \nabla_1 * \dots * \nabla_r$, and the projection along $\text{aff}(v_1, \dots, v_{r'})$ yields $\nabla_{I_1} * \nabla_{r'+1} * \dots * \nabla_r$ as the Gorenstein polytope dual to $\Delta^{I_1} * \Delta_{r'+1} * \dots * \Delta_r$. Hence, eventually we get $\nabla_{I_1} * \nabla_{I_2} * \dots * \nabla_{I_l}$ as the Gorenstein polytope of index l dual $\Delta^{I_1} * \Delta^{I_2} * \dots * \Delta^{I_l}$.

6.2. Projecting. Here, we show that projecting along a lower-dimensional lattice polytope in a nef-partition still yields a nef-partition.

The following definition is convenient:

Definition 6.5. Let $A \subseteq M_{\mathbb{R}}$ be a set. Then

$$A^\perp := \text{lin}(A)^\perp \subseteq N_{\mathbb{R}}.$$

We start with a general observation:

Lemma 6.6. *Let P, Q be lattice polytopes with $\dim(P) < \dim(P + Q)$. We denote by π the projection along $\text{aff}(P)$. Then preimages of facets of $\pi(P + Q)$ are facets of $P + Q$.*

Proof. By translation we may assume $\text{lin}(P) = \text{aff}(P)$. Let G be a facet of $\pi(P + Q)$. Then the preimage F of G is a Minkowski sum $F = P + F_Q$ for a face F_Q of Q . Since $\pi(P + Q) = \pi(Q)$, we easily see that $\dim(F) = \dim(P) + \dim(G)$. Now, the statement follows from $\dim(G) = \dim(P + Q) - \dim(P) - 1$. $\square$

Corollary 6.7. *Let P, Q be lattice polytopes with $P + Q$ reflexive. Then the projection $\pi(P + Q)$ along $\text{aff}(P)$ is a reflexive polytope.*

Proof. We may suppose $\dim(P) < \dim(P + Q)$. Let m be the unique interior lattice point of $P + Q$. By considering $P - m, Q$ instead of P, Q we may assume that $P + Q$ is reflexive with respect to 0. Let p be any lattice point of P . Then by considering $P - p, Q + p$ instead of P, Q we may also assume that $0 \in P$. Hence $\text{aff}(P) = \text{lin}(P)$. So $\pi(P + Q)$ has 0 as an interior lattice point. Now, Lemma 6.6 yields that any vertex of $(\pi(P + Q))^* = (P + Q)^* \cap P^\perp$ is also a vertex of $(P + Q)^*$. Since $(P + Q)^*$ is a lattice polytope, $(\pi(P + Q))^*$ is also a lattice polytope. $\square$

Now, we deal with our special situation of dual to each other centered nef-partitions $\Delta_1, \dots, \Delta_r$ and $\nabla_1, \dots, \nabla_r$:

Proposition 6.8. *Let $\{1, \dots, r\} = I \sqcup J$ with $I \neq \emptyset \neq J$, such that $\text{lin}(\Delta^J)$ has dimension strictly lower than $M_{\mathbb{R}}$. We denote by π the projection map along $\text{lin}(\Delta^J)$. Then $\pi(\Delta_1), \dots, \pi(\Delta_r)$ is a centered nef-partition.*

The dual reflexive polytope (in $(\Delta^J)^\perp$)

$$(\pi(\Delta_1) + \dots + \pi(\Delta_r))^* = \nabla_I \cap (\Delta^J)^\perp$$

is the smallest face F of ∇_I containing 0.

The dual nef-partition is given as $\nabla_1 \cap F, \dots, \nabla_r \cap F$.

Proof. By Cor. 6.3, we have

$$(\mathbb{R}_{\geq 0}(\Delta^J))^\vee = \mathbb{R}_{\geq 0}(\nabla_I),$$

and

$$\begin{aligned} (\pi(\Delta_1) + \dots + \pi(\Delta_r))^* &= (\Delta_1 + \dots + \Delta_r)^* \cap (\Delta^J)^\perp = \\ &= \text{Conv}(\nabla_I, \nabla_J) \cap (\Delta^J)^\perp = \nabla_I \cap (\Delta^J)^\perp = F. \end{aligned}$$

The other statements are clear. $\square$

Corollary 6.9. *Let $\{1, \dots, r\} = I \sqcup J$ with $I \neq \emptyset \neq J$. Let F be the smallest face F of ∇_I containing 0. Then*

$$\dim(F) + \dim(\Delta^J) = \dim(M_{\mathbb{R}}).$$

In particular, $\dim(\Delta_1) = \dim(M_{\mathbb{R}})$ iff 0 is a vertex of $\text{Conv}(\nabla_2, \dots, \nabla_r)$.

6.3. Decomposing. In [BB96a] Borisov and the first author introduced the notion of an irreducible nef-partition:

Definition 6.10. Let $\emptyset \neq I \subseteq \{1, \dots, r\}$ such that Δ^I is a reflexive polytope (in its linear span). By Prop. 6.8 it is enough to assume that Δ^I contains 0 in its relative interior. We say I is *irreducible*, if I is minimal with this property.

The proper nef-partition $\Delta_1, \dots, \Delta_r$ is called *irreducible*, if $\{1, \dots, r\}$ is irreducible.

Theorem 5.8 of [BB96a] states that any (proper) nef-partition has a unique *decomposition* into irreducible (proper) nef-partitions, i.e., there are unique irreducible subsets $I_1, \dots, I_l$ that form a partition of $\{1, \dots, r\}$ such that

$$\Delta_1 + \dots + \Delta_r = \Delta^{I_1} + \dots + \Delta^{I_l}, \quad \text{lin}(\Delta^{I_1}) \oplus \dots \oplus \text{lin}(\Delta^{I_l}) = M_{\mathbb{R}}.$$

This is called *semi-simplicity of nef-partitions*. Here, we recall the proof of the existence of such a decomposition and give a generalization in Prop. 6.13. We also prove in Prop. 6.15 the uniqueness of this decomposition, which was missing in [BB96a]. Finally, we determine the maximal length of a proper nef-partition in Prop. 6.16.

The existence of a decomposition into irreducible nef-partitions is an immediate consequence of the following result, which was shown in [BB96a, Thm.5.8] (we include a proof for the convenience of the reader):

Proposition 6.11. *Let $\{1, \dots, r\} = I \sqcup J$ with $I \neq \emptyset \neq J$. If Δ^I is reflexive, then also Δ^J is reflexive, and $\text{lin}(\Delta^I) \oplus_{\mathbb{R}} \text{lin}(\Delta^J) = M_{\mathbb{R}}$.*

Proof. By Prop. 6.2 Δ^I, Δ^J and ∇_I, ∇_J are dual nef-partitions. Since Δ^I contains 0 in its relative interior, Prop. 6.8 yields $\Delta^I \subseteq U := \nabla_J^\perp$. Moreover, $\Delta^J \subseteq \sigma := (\mathbb{R}_{\geq 0}(\nabla_I))^\vee$ because of $\langle \Delta^J, \nabla_I \rangle \geq 0$. Furthermore, $U \cap \sigma = \{0\}$, since $\langle x, \nabla_I + \nabla_J \rangle \geq 0$ is only possibly for $x = 0$. We show that σ is actually a linear subspace of $M_{\mathbb{R}}$. This already yields $U \oplus \sigma = \text{lin}(\Delta^I) \oplus \text{lin}(\Delta^J) = M_{\mathbb{R}}$, and by projecting along $U = \text{lin}(\Delta^I)$ we get that Δ^J is reflexive by Prop. 6.8.

Let $x \in \sigma$. We have to show $-x \in \sigma$. Let ρ be the projection along U . As 0 is in the interior of $\Delta^I + \Delta^J$, and $\rho(\Delta^I) = \{0\}$, we get $\rho(\sigma) \supseteq \rho(\mathbb{R}_{\geq 0}(\Delta^J)) = \rho(M_{\mathbb{R}})$. Therefore there is some $x' \in \sigma$ with $\rho(x') = \rho(-x)$. This implies $x' + x \in U \cap \sigma = \{0\}$, thus $-x = x' \in \sigma$. $\square$

As was observed in [BB96a, Example 5.5], note that the direct sum in Prop. 6.11 is in general only a splitting of vector spaces, *not* of lattices.

Corollary 6.12. *Let $\{1, \dots, r\} = I \sqcup J$ with $I \neq \emptyset \neq J$. If one of the lattice polytopes in $\{\Delta^I, \Delta_I, \Delta^J, \Delta_J, \nabla^I, \nabla_I, \nabla^J, \nabla_J\}$ contains 0 in its relative interior, then anyone does, and, moreover, each one is a reflexive polytope.*

In this case: $(\Delta_i)_{i \in I}$ is a nef-partition with dual nef-partition $(\nabla_i)_{i \in I}$, with respect to the dual vector spaces $\nabla_J^\perp \subseteq M_{\mathbb{R}}$ and $\Delta_J^\perp \subseteq N_{\mathbb{R}}$.

Proof. By Prop. 6.2, Δ^I, Δ^J and ∇_I, ∇_J are dual nef-partitions. The same is true for ∇^I, ∇^J and Δ_I, Δ_J . Hence, Prop. 6.8, implies that if some of these 8 polytopes contains 0 in its relative interior, then it is necessarily a reflexive polytope. Since Δ^I has 0 in its relative interior if and only if Δ_I has 0 in its relative interior, we may assume that Δ^I is reflexive. Prop. 6.11 yields that also Δ^J is reflexive. Now, Cor. 6.3 implies

$$\nabla_I = \{y \in (\mathbb{R}_{\geq 0}(\Delta^J))^\vee : \langle \Delta^I, y \rangle \geq -1\} = \{y \in (\Delta_J)^\perp : \langle \Delta^I, y \rangle \geq -1\},$$

since Δ_J has 0 in its relative interior. From this, we get $\nabla_I = (\Delta^I)^*$. Hence, ∇_I is reflexive. In the same way we see $\nabla_J = (\Delta^J)^*$, so also ∇_J is reflexive. Now, the last statement follows merely from the definition of dual nef-partitions. $\square$

There is the following generalization of Prop. 6.11 (this result is joint work with Christian Haase):

Proposition 6.13. *Let P, Q be lattice polytopes with $P + Q$ reflexive. If P contains a lattice point in its relative interior, then P and Q are reflexive polytopes, and P, Q form a nef-partition.*

Proof. As at the beginning of the proof of Cor. 6.7 we may assume that 0 is an interior lattice point of P and $P + Q$.

We define $Q' := Q \cap \text{lin}(P)$. Since $0 \in P + Q$, there exist $x \in P, y \in Q$ with $x + y = 0$. Hence $Q' \neq \emptyset$. Assume there is some $q \in Q', q \neq 0$. Since 0 is in the relative interior of P , there is a linear form u in the dual space of $\text{lin}(P)$ such that $\langle u, q \rangle < 0$, and $F := \{x \in P : \langle u, x \rangle = -1\}$ is a facet of P . Let $y \in Q'$ with $\langle u, y \rangle = \min_{x \in Q'} \langle u, x \rangle =: c$. We have $c \leq \langle u, q \rangle < 0$. Hence, $G := \{x \in P + Q' : \langle u, x \rangle = -1 + c\}$ is a face of $P + Q' = (P + Q) \cap \text{lin}(P)$. Since $0 \notin G$, $F + y \subseteq G$ and $\dim(F) = \dim(\text{lin}(P)) - 1$, we see that G is a facet of $(P + Q) \cap \text{lin}(P)$, parallel to F . Since $P + Q$ is reflexive, there is an integral linearform w in the dual space of $\text{lin}(P)$ such that $\langle w, G \rangle = -1$. We have $u = \lambda w$ for $\lambda := 1 - c > 1$. Let v be a vertex of F , hence $\langle w, v \rangle \in \mathbb{Z}$. This implies $\lambda = \langle u, v \rangle / \langle w, v \rangle = -1 / \langle w, v \rangle \leq 1$, a contradiction.

We have shown that $Q \cap \text{lin}(P) = \{0\}$. In particular $0 \in Q$, thus P, Q is by definition a centered nef-partition. Now, apply Cor. 6.12 and Prop. 6.11. $\square$

Example 6.14. Let $P := \text{Conv}((1, 0, 0), (0, 1, 0), (-1, -1, 0))$ and $Q := \text{Conv}((1, 0, -1), (0, 1, 1))$. Then $P + Q$ has only one interior lattice point, P is reflexive, however Q has no interior lattice point. This example, due to Christian Haase, shows that in Prop. 6.13 the assumption that $P + Q$ is reflexive cannot be weakened by the condition that $P + Q$ has only one interior lattice point.

The uniqueness of a decomposition into irreducible nef-partitions is a corollary of the following result:

Proposition 6.15. *Any two irreducible subsets of $\{1, \dots, r\}$ are disjoint.*

Proof. Assume there exist irreducible subsets I, I' of $\{1, \dots, r\}$ with $I \neq I'$ and $A := I \cap I' \neq \emptyset$. Thus, we have the following situation: $\{1, \dots, r\} = I \sqcup J = I' \sqcup J'$, $I = A \sqcup B$, $J = C \sqcup D$, $I' = A \sqcup C$, $J' = B \sqcup D$.

By Prop. 6.11 we have $\text{lin}(\Delta^I) \cap \text{lin}(\Delta^{J'}) = \{0\}$, thus $\text{lin}(\Delta^A) \cap \text{lin}(\Delta^C) = \{0\}$. Hence, since $\Delta^{I'} = \Delta^A + \Delta^C$ contains 0 in its relative interior, also Δ^A contains 0 in its relative interior. This contradicts I being irreducible. $\square$

Finally, we examine nef-partitions of largest length:

Proposition 6.16. *A proper nef-partition has length $r \leq 2 \dim(M_{\mathbb{R}})$, with equality only if the convex hull is combinatorially a crosspolytope, i.e., the dual of a cube.*

Proof. An irreducible proper nef-partition has length $r \leq \dim(M_{\mathbb{R}}) + 1$, see [BB96a, Cor.3.5] or [HZ05, Lemma 2.6]. Now, a proper nef-partition of length r decomposes into, say l , irreducible nef-partitions. This yields

$$r \leq \dim(M_{\mathbb{R}}) + l \leq 2 \dim(M_{\mathbb{R}}),$$

where complete equality holds if and only if any irreducible nef-partition has dimension one and length two. $\square$

REFERENCES

- [Ath05] C.A. Athanasiadis, *Ehrhart polynomials, simplicial polytopes, magic squares and a conjecture of Stanley*, J. Reine Ang. Math. **583** (2005), 163–174.
- [Bat82] V.V. Batyrev, *Boundedness of the degree of multidimensional toric Fano varieties*, Mosc. Univ. Math. Bull. **37** (1982), 28–33.
- [Bat94] V.V. Batyrev, *Dual polyhedra and mirror symmetry for Calabi-Yau hypersurfaces in toric varieties*, J. Algebr. Geom. **3** (1994), 493–535.
- [BS95] V.V. Batyrev, D. van Straten, *Generalized hypergeometric functions and rational curves on Calabi-Yau complete intersections in toric varieties*, Comm. Math. Phys. **168** (1995), no. 3, 493–533.
- [Bat98] V.V. Batyrev, *Stringy Hodge numbers of varieties with Gorenstein canonical singularities*, In: M.-H. Saito (ed.) et al., Integrable systems and algebraic geometry, Proceedings of the 41st Taniguchi symposium, Japan 1997, Singapore: World Scientific, 1–32, 1998.
- [Bat00] V.V. Batyrev, *Stringy Hodge numbers and Virasoro algebra*, Math. Res. Lett. **7** (2000), 155–164.
- [Bat07] V.V. Batyrev, *Lattice polytopes with a given h^* -polynomial*, In: C.A. Athanasiadis (ed.) et al., Algebraic and geometric combinatorics, Proceedings of a Euroconference in Mathematics, Anogia, Crete, Greece, August 20–26, 2005, Providence, RI: AMS, Contemp. Math. **423**, 1–10, 2007.
- [BB96a] V.V. Batyrev & L.A. Borisov, *On Calabi-Yau complete intersections in toric varieties*, In: M. Andreatta (ed.) et al., Higher dimensional complex varieties, Proceedings of the international conference, Trento, Italy, June 15–24, 1994, Berlin: Walter de Gruyter, 39–65, 1996.
- [BB96b] V.V. Batyrev & L.A. Borisov, *Mirror duality and string-theoretic Hodge numbers*, Invent. Math. **126** (1996), 183–203.
- [BB97] V.V. Batyrev & L.A. Borisov, *Dual cones and mirror symmetry for generalized Calabi-Yau manifolds*, In: B. Greene (ed.) et al., Mirror symmetry II,

- Cambridge, MA: International Press, AMS/IP Stud. Adv. Math. **1**, 71–86, 1997.
- [BD96] V.V. Batyrev & D.I. Dais, *Strong McKay correspondence, string-theoretic Hodge numbers and mirror symmetry*, Topology **35** (1996), 901–929.
- [BN06] V.V. Batyrev & B. Nill, *Multiples of lattice polytopes without interior lattice points*, Preprint, math.CO/0602336, 2006.
- [BorBor93] A. A. Borisov & L.A. Borisov, *Singular toric Fano varieties*, Math. USSR, Sb. **75** (1993), 277–283.
- [Bor93] L.A. Borisov, *Towards the mirror symmetry for Calabi-Yau complete intersections in Gorenstein toric Fano varieties*, Preprint, arxiv.AG/9310001, 1993.
- [BM03] L.A. Borisov & A.R. Mavlyutov, *String cohomology of Calabi-Yau hypersurfaces via mirror symmetry. Appendix: G-polynomials*, Adv. Math. **180** (2003), 355–390.
- [Bra03] T. Braden, *Koszul duality for toric varieties*, Preprint, math.AG/0308216, 2003.
- [Bra05] T. Braden, *Remarks on the combinatorial intersection cohomology of fans*, Preprint, math.CO/0511488, 2005.
- [BR05] W. Bruns & T. Roemer, *h-vectors of Gorenstein polytopes*, Preprint, math.AC/0508392, 2005.
- [CDP93] P. Candelas, E. Derrick, L. Parkes, *Generalized Calabi-Yau Manifolds and the Mirror of a Rigid Manifold*, Nucl. Phys. **B407** (1993), 115–154.
- [ET05] I.Z. Emiris & E.P. Tsigaridas, *Minkowski decomposition of convex lattice polygons*, In: M. Elkadi, B. Mourrain, R. Piene (eds.), Algebraic geometry and geometric modeling, Berlin: Springer-Verlag, 2005.
- [Gro05] M. Gross, *Toric degenerations and Batyrev-Borisov duality*, Math. Ann. **333** (2005), 645–688.
- [Haa00] C. Haase, *Lattice Polytopes and Unimodular Triangulations*, PhD-thesis, Technical University of Berlin, 2000.
- [Haa05] C. Haase, *Reflexive polytopes in dimension 2 and 3, and the numbers 12 and 24*, In: M. Beck, B. Chen, L. Fukshansky, C. Haase, A. Knutson, B. Reznick, S. Robins, and A. Schürmann, Problems from the Cottonwood Room, Contemporary Mathematics **374** (2005), 179–191.
- [HZ05] C. Haase & I. Zharkov, *Integral affine structures on spheres III: complete intersections*, Preprint, math.AG/0504181, 2005.
- [Hib92] T. Hibi, *Dual polytopes of rational convex polytopes*, Combinatorica **12** (1992), 237–240.
- [Ka05] Y. Kawamata, *Log crepant birational maps and derived categories*, J. Math. Sci. Univ. Tokyo **12** (2005), 211–231.
- [KKRS05] A. Klemm, M. Kreuzer, E. Riegler, E. Scheidegger, *Topological String Amplitudes, Complete Intersection Calabi-Yau Spaces and Threshold Corrections*, JHEP**05** (2005), 023.
- [KRS03] M. Kreuzer, E. Riegler, D. Sahakyan, *Toric complete intersections and weighted projective space*, J.Geom.Phys. **46** (2003), 159–173.
- [KS98] M. Kreuzer & H. Skarke, *Classification of reflexive polyhedra in three dimensions*, Adv. Theor. Math. Phys. **2** (1998), 853–871.
- [KS00] M. Kreuzer & H. Skarke, *Complete classification of reflexive polyhedra in four dimensions*, Adv. Theor. Math. Phys. **4** (2000), 1209–1230.
- [KS04] M. Kreuzer & H. Skarke, *PALP, A package for analyzing lattice polytopes with applications to toric geometry*, Computer Phys. Comm. **157** (2004), 87–106.
- [NH97] E De Nigri & T. Hibi, *Gorenstein algebras of Veronese type*, J. Algebra **193** (1997), 629–639.

- [OH05] H. Ohsugi & T. Hibi, *The h -vector of a Gorenstein toric ring of a compressed polytope*, Electr. J. Comb. **11** (2005), Research paper N4.
- [RW05] V. Reiner & V. Welker, *On the Charney-Davis and Neggers-Stanley conjectures*, J. Comb. Th. A **109** (2005), 247–280.
- [Sta87] R. Stanley, *Generalized h -vectors, intersection cohomology of toric varieties, and related results*, In: Commutative algebra and combinatorics, US-Jap. Joint Semin., Kyoto/Jap. 1985, Adv. Stud. Pure Math. **11** (1987), 187–213.
- [Sta92] R. Stanley, *Subdivisions and local h -vectors*, J. Am. Math. Soc. **5** (1992), 805–851.
- [Sti98] J. Stienstra, *Resonant hypergeometric systems and mirror symmetry*, In: M.-H. Saito (ed.) et al., Integrable systems and algebraic geometry, Proceedings of the 41st Taniguchi symposium, Kobe, Japan, June 30–July 4, 1997, and in Kyoto, Japan, July 7–11, 1997, Singapore: World Scientific, 412–452, 1998.
- [Va89] C. Vafa, *String vacua and orbifoldized LG-models*, Mod. Phys. Lett. **A4** (1989), 1169.

MATHEMATISCHES INSTITUT, UNIVERSITÄT TÜBINGEN, AUF DER MORGENSTELLE 10,
72076 TÜBINGEN, GERMANY

E-mail address: `victor.batyrev@uni-tuebingen.de`

ARBEITSGRUPPE GITTERPOLYTOPE, FREIE UNIVERSITÄT BERLIN, ARNIMALLEE 3,
14195 BERLIN, GERMANY

E-mail address: `nill@math.fu-berlin.de`